

THREEFOLD EXTREMAL CONTRACTIONS OF TYPE (IA)

SHIGEFUMI MORI AND YURI PROKHOROV

Dedicated to the memory of Professor Masaki Maruyama

ABSTRACT. Let (X, C) be a germ of a threefold X with terminal singularities along an irreducible reduced complete curve C with a contraction $f : (X, C) \rightarrow (Z, o)$ such that $C = f^{-1}(o)_{\text{red}}$ and $-K_X$ is ample. Assume that a general member $F \in |-K_X|$ meets C only at one point P and furthermore (F, P) is Du Val of type A if $\text{index}(X, P) = 4$. We classify all such germs in terms of a general member $H \in |\mathcal{O}_X|$ containing C .

CONTENTS

1. Introduction	1
2. Preliminaries	5
3. Deformations of 3-dimensional divisorial contractions	9
4. Case cD/3	14
5. Case: P is of type cA/m and H is normal	25
6. Case: P is of type cA/m and H is not normal	27
7. On index two $\mathbb{Q}$ -conic bundles	35
References	36

1. INTRODUCTION

1.1. Definition. Let (X, C) be a germ of a threefold with terminal singularities along an reduced complete curve. We say that (X, C) is an *extremal curve germ* if there is a contraction $f : (X, C) \rightarrow (Z, o)$ such that $C = f^{-1}(o)_{\text{red}}$ and $-K_X$ is f -ample.

If furthermore f is birational, then (X, C) is said to be an *extremal neighborhood* [Mor88]. In this case f is called *flipping* if its exceptional locus coincides with C (and then (X, C) is called *isolated*). Otherwise the exceptional locus of f is two-dimensional and f is called *divisorial*. If f is not birational, then $\dim Z = 2$ and (X, C) is said to be a $\mathbb{Q}$ -conic bundle germ [MP08].

In this paper, unless explicitly stated otherwise, we assume that C is *irreducible*.

1.2. Let (X, C) be an extremal curve germ as above. For each singular point P of X with $P \in C$, consider the germ $(P \in C \subset X)$. All such germs (or all such singular points, for simplicity) are classified into types: (IA), (IC), (IIA), (IIB), (III), (IA $^\vee$), (II $^\vee$), (ID $^\vee$), (IE $^\vee$), as for whose definitions we refer the reader to [Mor88] and [MP08]. The possible configurations of such points are also classified in [Mor88] and [MP08]. Moreover, it is known that a general member $F \in |-K_X|$ has

only Du Val singularities and all possibilities for F are described [Mor88, 7.3, 9.10], [KM92, 2.2], [MP08, 1.3.7], [MP09, 2.1-2.2]. The next step in the classification is to study a general hyperplane section, that is, a general divisor H of $|\mathcal{O}_X|_C$, the linear subsystem of $|\mathcal{O}_X|$ consisting of sections $\supset C$. Roughly speaking, the importance of this divisor can be explained as follows. Once we have this H , the total threefold can be considered as a one-parameter deformation of H . Then one can apply the deformation theory to construct X starting from a two-dimensional data $H \supset C$.

In this paper we classify extremal curve germs of type (IA) or $(\text{IA}^\vee)$ in terms of a general member $H \in |\mathcal{O}_X|_C$. An extremal curve germ (X, C) is said to be of type (IA) (resp. $(\text{IA}^\vee)$) if it contains exactly one non-Gorenstein point P and it is of type (IA) (resp. $(\text{IA}^\vee)$). For readers' convenience, we note the following characterization (cf. [KM92, Th. 2.2]) for an extremal curve germ (X, C) with a point P of index $m > 1$ to be of type (IA) or $(\text{IA}^\vee)$ in terms of a general member $F \in |-K_X|$: (X, C) is of type (IA) or $(\text{IA}^\vee)$ if and only if (i) $F \cap C = \{P\}$ as a set, and (ii) (F, P) is Du Val of type A if $m = 4$.

1.3. Throughout this paper if we do not specify otherwise we assume that (X, C) of type (IA) or $(\text{IA}^\vee)$. More precisely, X contains a unique non-Gorenstein terminal point $P \in X$, which is of type (IA) or $(\text{IA}^\vee)$.

A point $(X \supset C \ni P)$ of index $m > 1$ is said to be of type (IA) if there exists an embedding $X \subset \mathbb{C}_{x_1, \dots, x_4}^4 / \mu_m(a_1, a_2, -a_1, 0)$ such that

$$C = \{x_1^{a_2} - x_2^{a_1} = x_3 = x_4 = 0\} / \mu_m(a_1, a_2, -a_1, 0),$$

for some positive integers a_1, a_2 with $\gcd(a_1 a_2, m) = 1$ and $m \in a_1 \mathbb{Z}_{>0} + a_2 \mathbb{Z}_{>0}$, and X is given by an invariant vanishing along C [Mor88, A.3]. If f is a $\mathbb{Q}$ -conic bundle, then $a_2 = 1$ by [MP08, Proposition 8.5]. Points of type $(\text{IA}^\vee)$ are described similarly [Mor88, A.3].

For a normal surface S and a curve $V \subset S$, we use the usual notation of graphs $\Delta(S, V)$ of the minimal resolution of S near V : each $\diamond$ corresponds to an irreducible component of V and each $\circ$ corresponds to an exceptional divisor on the minimal resolution of S , and we may use $\bullet$ instead of $\diamond$ if we want to emphasize that it is a complete (-1) -curve. A number attached to a vertex denotes the minus self-intersection number. For short, we may omit 2 if the self-intersection is -2 .


Flipping extremal neighborhoods containing a terminal singular point of type cD/3 ([Mor85], [Rei87]) are classified in [KM92, Theorems 6.2 and 6.3]. Thus the following theorem covers the rest of extremal curve germs which contain cD/3 points.

1.4. Theorem. *Let (X, C) be an extremal curve germ. Assume that (X, C) is of type (IA) and let $P \in X$ be the non-Gorenstein point. Assume furthermore that (X, P) is of type cD/3. Then f is a birational contraction, not a $\mathbb{Q}$ -conic bundle. The general member $H \in |\mathcal{O}_X|_C$ and its image $T = f(H) \in |\mathcal{O}_Z|$ are normal and have only rational singularities. Moreover, if f is not a flipping contraction, then the following are the only possibilities for the dual graphs of (H, C) and T :*

$$1.4.1. \quad \Delta(H, C): \quad \circ - \bullet - \overset{3}{\circ} - \underset{\overset{3}{\circ}}{\underset{\underset{3}{\circ}}{\circ}} - \overset{3}{\circ}$$


and T is of type A_2 ; here (X, P) is a simple cD/3 point (see 4.1);

1.4.2. $\Delta(H, C)$:



and T is of type D_4 ; here (X, P) is a double cD/3 point.

1.4.3. $\Delta(H, C)$:



and T is of type E_6 ; here (X, P) is a triple cD/3 point.

In all the cases above the right hand side of the graph for (H, C) corresponds to the non-Gorenstein point $P \in H$. The left hand side corresponds to either a type (III) point or a smooth point of X .

Examples are given in 4.14.1 and 4.14.2.

Note that $\mathbb{Q}$ -conic bundles of type $(IA^\vee)$ are completely classified in [MP08]. The following two theorems cover the $\mathbb{Q}$ -conic bundles of type (IA).

1.5. Theorem. *Let $(X, C \simeq \mathbb{P}^1)$ be a $\mathbb{Q}$ -conic bundle germ of index $m > 2$ and of type (IA). Let $P \in X$ be the non-Gorenstein point. Then (X, P) is a point of type cA/ m and a general member $H \in |\mathcal{O}_X|_C$ is not normal. Furthermore, the dual graph of (H', C') , the normalization H' and the inverse image C' of C is of the form*

$$\underbrace{\overset{a_r}{\circ} \cdots \overset{a_1}{\circ}}_{\Delta_1} - \bullet - \underbrace{\overset{b_1}{\circ} \cdots \overset{b_s}{\circ}}_{\Delta_2}$$

(in particular, C' is irreducible). Here the chain Δ_1 (resp., Δ_2) corresponds to the singularity of type $\frac{1}{m}(1, a)$ (resp., $\frac{1}{m}(1, -a)$) for some integer $a \in [1, m]$ relatively prime to m . The germ (H, C) is analytically isomorphic to the germ along the line $y = z = 0$ of the hypersurface given by the following weighted polynomial of degree $2m$ in variables x, y, z, u :

$$\phi := x^{2m-2a}y^2 + x^{2a}z^2 + yzu.$$

in $\mathbb{P}(1, a, m-a, m)$. Furthermore (X, C) is given as an analytic germ of a subvariety of $\mathbb{P}(1, a, m-a, m) \times \mathbb{C}_t$ along $C \times 0$ given by

$$\phi + \alpha_1 x^{2m-a}y + \alpha_2 x^{m-a}uy + \alpha_3 x^{2m} + \alpha_4 x^m u + \alpha_5 u^2 = 0$$

for some $\alpha_1, \dots, \alpha_5 \in t\mathcal{O}_{0, \mathbb{C}_t}$ and there is a $\mathbb{Q}$ -conic bundle structure $X \rightarrow \mathbb{C}^2$ through which the second projection $X \rightarrow \mathbb{C}_t$ factors.

An explicit example is given in 6.8.4.

1.6. Theorem ([Pro97, §3], [MP08, Th. 12.1]). *Let $(X, C \simeq \mathbb{P}^1)$ be a $\mathbb{Q}$ -conic bundle germ of index 2 and of type (IA). Let $f: (X, C) \rightarrow (Z, o)$ be the corresponding contraction. Then (Z, o) is smooth. Let u, v be a local coordinates on (Z, o) . Then there is an embedding*

$$f: X \hookrightarrow \mathbb{P}(1, 1, 1, 2) \times Z \xrightarrow{p} Z$$

such that X is given by two equations

$$\begin{aligned} q_1(y_1, y_2, y_3) &= \psi_1(y_1, \dots, y_4; u, v), \\ q_2(y_1, y_2, y_3) &= \psi_2(y_1, \dots, y_4; u, v), \end{aligned}$$

where ψ_i and q_i are weighted quadratic in $y_1, \dots, y_4$ with respect to $\text{wt}(y_1, \dots, y_4) = (1, 1, 1, 2)$ and $\psi_i(y_1, \dots, y_4; 0, 0) = 0$. The only non-Gorenstein point of X is $(0, 0, 0, 1; 0, 0)$. Up to projective transformations, the following are the only possibilities for q_1 and q_2 :

- (i) $q_1 = y_1^2$, $q_2 = y_2^2 - y_1 y_3$; here a general member $H \in |\mathcal{O}_X|_C$ is normal.
- (ii) $q_1 = y_1^2$, $q_2 = y_2^2$; here every member $H \in |\mathcal{O}_X|_C$ is non-normal.

In both cases C is given by $u = v = y_1 = y_2 = 0$.

Explicit examples are given in Sect. 7 (see also Remark 6.7.1).

1.7. For a triple (X, C, P) of type (IA) or $(\text{IA}^\vee)$, the singularity (X, P) is either cA/m, cD/3 or of index 2. Extremal neighborhoods of index 2 are classified in [KM92, §4]. The cD/3 case is covered by [KM92, §6] and Theorem 1.4. The next theorem covers the remaining case.

1.8. Theorem (cf. [Tzi05]). *Let (X, C) be an extremal neighborhood of type (IA) or $(\text{IA}^\vee)$. Let $P \in X$ be the non-Gorenstein point. Assume furthermore that (X, P) is of type cA/m. Let $F \in |-K_X|$ be a general member. Then there exists a member $H \in |\mathcal{O}_X|_C$ such that the pair $(X, H + F)$ is LC.*

1.8.1. *If H is normal, then H has only log terminal singularities of type T. The graph $\Delta(H, C)$ is of the form*

$$(1.8.1.1) \quad \begin{array}{c} \circ^{c_1} - \circ^{c_2} - \dots - \circ^{c_r} - \dots - \circ^{c_n} \\ | \\ \bullet \\ | \\ \circ - \circ - \dots - \circ \end{array}$$

Here the chain $[c_1, \dots, c_n]$ corresponds to the non-Du Val singularity (H, P) of type T. The chain of (-2) -vertices in the last line corresponds to a Du Val point (H, Q) . It is possible that this chain is empty (i.e., (H, Q) is smooth). Cases $r = 1$ and $r = n$ are also not excluded.

1.8.2. *If every member of $|\mathcal{O}_X|_C$ is non-normal, then the dual graph of the normalization (H', C') is of the form*

$$(1.8.2.1) \quad \underbrace{\circ^{a_r} - \dots - \circ^{a_1}}_{\Delta_1} - \bullet - \underbrace{\circ^{c_1} - \dots - \circ^{c_l}}_{\Delta_3} - \diamond - \underbrace{\circ^{b_1} - \dots - \circ^{b_s}}_{\Delta_2}$$

(in particular, C' is reducible). The chain Δ_1 (resp., Δ_2) corresponds to the singularity of type $\frac{1}{m}(1, a)$ (resp., $\frac{1}{m}(1, -a)$) for some a with $\gcd(m, a) = 1$ and the chain Δ_3 corresponds to the point (H', Q') , where $Q' = C'_1 \cap C'_2$. The strings $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate (cf. 2.1.2). Moreover,

$$\sum (c_i - 2) \leq 2 \quad \text{and} \quad \tilde{C}_1^2 + \tilde{C}_2^2 + 5 - \sum (c_i - 2) \geq 0,$$

where $\tilde{C} = \tilde{C}_1 + \tilde{C}_2$ is the proper transform of C on the minimal resolution $\tilde{H}$. Both components of $\tilde{C}$ are contracted on the minimal model of $\tilde{H}$. In this case, the triple (X, C, P) is analytically isomorphic to $(\{\alpha = 0\}, x_1\text{-axis}, 0)/\mu_m(1, a, -a, 0)$, where $\gcd(m, a) = 1$ and $\alpha(x_1, \dots, x_4) = 0$ is the equation of a terminal cA/m-point in $\mathbb{C}^4/\mu_m(1, a, -a, 0)$. (In particular, (X, C) is of type (IA)).

Conversely, for any germ $(H, C \simeq \mathbb{P}^1)$ of the form 1.8.1 or 1.8.2 admitting a birational contraction $(H, C) \rightarrow (T, o)$ there exists a threefold birational contraction $f : (X, C) \rightarrow (Z, o)$ as in 1.1 of type (IA) such that $H \in |\mathcal{O}_X|_C$.

1.8.3. Remark. Basically this result is proved in [Tzi05]. However [Tzi05] treated only *divisorial* contractions that contracts a divisor to a *smooth* curve. Under these assumptions the result of [Tzi05] is much stronger.

1.8.4. Remark. Note that in 1.8 H is not assumed to be a general element of $|\mathcal{O}_X|_C$. If H is chosen general, then the cases 1.8.1 and 1.8.2 cover all the cases under 1.8. Proposition 6.3 gives a criterion for a general member of $|\mathcal{O}_X|_C$ to be non-normal, and Proposition 6.6 gives, under some additional assumptions, a criterion for a given H to be general.

To check divisoriality one can use the following criterion which is an immediate consequence of Theorem 3.1.

1.9. Theorem. *Let $f : (X, C \simeq \mathbb{P}^1) \rightarrow (Z, o)$ be a 3-dimensional birational extremal curve germ. Then f is divisorial if and only if (Z, o) is a terminal singularity.*

One of our technical tools is the deformation of extremal curve germs. In particular, the following result shows that for every extremal curve germ $f : (X, C) \rightarrow (Z, o)$ the contraction f deforms with X . Combined with Theorem 1.9, it allows us to run MMP for every deformation of an extremal curve germ which may not be $\mathbb{Q}$ -factorial.

1.10. Theorem (cf. [KM92, (11.4)], [MP08, (6.2)]). *Let $f : (X, C) \rightarrow (Z, o)$ be an extremal divisorial (resp. flipping, $\mathbb{Q}$ -conic bundle) curve germ, where C is not necessarily irreducible. Let $\pi : \mathcal{X} \rightarrow (\mathbb{C}_\lambda^1, 0)$ be a flat deformation of $X = \mathcal{X}_0 := \pi^{-1}(0)$ over a germ $(\mathbb{C}_\lambda^1, 0)$ with a flat closed subspace $\mathcal{C} \subset \mathcal{X}$ such that $\mathcal{C} = \mathcal{C}_0$. Then there exist a flat deformation $\mathcal{Z} \rightarrow (\mathbb{C}_\lambda^1, 0)$ and a proper $\mathbb{C}_\lambda^1$ -morphism $\mathfrak{f} : \mathcal{X} \rightarrow \mathcal{Z}$ such that $f = \mathfrak{f}_0$ and $\mathfrak{f}_\lambda : (\mathcal{X}_\lambda, \mathfrak{f}_\lambda^{-1}(o_\lambda)_{\text{red}}) \rightarrow (\mathcal{Z}_\lambda, o_\lambda)$ is a divisorial (resp. flipping, $\mathbb{Q}$ -conic bundle) extremal curve germ for every small λ , where $o_\lambda := \mathfrak{f}_\lambda(\mathcal{C}_\lambda)$.*

1.11. Conventions. We work over the complex number field $\mathbb{C}$. Notations and techniques of [Mor88] will be used freely. In particular, for a terminal singularity (X, P) the index-one cover is denoted by $(X^\sharp, P^\sharp) \rightarrow (X, P)$ and for a subvariety $V \subset X$ its preimage is denoted by $V^\sharp$.

2. PRELIMINARIES

2.1. Some facts about two-dimensional toric singularities.

2.1.1. Notation. A continued fraction

$$a_1 - \frac{1}{a_2 - \frac{1}{\ddots - \frac{1}{a_r}}} \quad (a_1, \dots, a_r \geq 2)$$

is denoted by $[a_1, \dots, a_r]$ and called a *string*. Write $m/q = [a_1, \dots, a_r]$, where $\gcd(m, q) = 1$. Given m and q this expression is unique. It is well-known that the minimal resolution of the cyclic quotient singularity $\frac{1}{m}(1, q)$ is a chain of smooth rational curves whose self-intersection numbers are $-a_1, \dots, -a_r$.

2.1.2. Definition. We say that a string $[b_1, \dots, b_s]$ is *conjugate* to $[a_1, \dots, a_r]$ if $[b_1, \dots, b_s] = m/(m - q)$.

2.1.3. Lemma.

- (i) If the strings $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate, then either $a_1 = 2$ or $b_1 = 2$.
- (ii) The strings $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ with $a_1 = 2$ and $r > 1$ are conjugate if and only if so are $[a_2, \dots, a_r]$ and $[b_1 - 1, \dots, b_s]$.
- (iii) The strings $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate if and only if so are $[a_r, \dots, a_1]$ and $[b_s, \dots, b_1]$.

2.2. T-singularities.

2.2.1. Definition ([KSB88]). A normal surface singularity is said to be of *type T* if it is log terminal and admits a $\mathbb{Q}$ -Gorenstein one-parameter smoothing.

2.2.2. Proposition ([LW86, Prop. 5.9], [KSB88, Prop. 3.10]). A surface singularity is of type T if and only if it is either Du Val or a cyclic quotient of type $\frac{1}{n}(a, b)$, where $\gcd(n, a) = \gcd(n, b) = 1$ and $(a + b)^2 \equiv 0 \pmod n$.

By 2.1 any non-Du Val T-singularity is represented by some string $[a_1, \dots, a_r]$. Then we say that $[a_1, \dots, a_r]$ is a *T-string*, or a *string of type T*.

2.2.3. Proposition ([KSB88, Prop. 3.11]).

- (i) The strings $[4]$, $[3, 3]$, and $[3, 2, \dots, 2, 3]$ are of type T.
- (ii) If the string $[a_1, \dots, a_r]$ is of type T, then so are $[2, a_1, \dots, a_{r-1}, a_r + 1]$ and $[a_1 + 1, a_2, \dots, a_r, 2]$.
- (iii) Every non-Du Val string of type T can be obtained by starting with one described in (i) and iterating the steps described in (ii).

2.2.4. Corollary. Let (X, P) be a $\mathbb{Q}$ -Gorenstein isolated threefold singularity and let $H \subset X$ be a surface such that H is a Cartier divisor. If the singularity (H, P) is log terminal, then (H, P) is a T-singularity and the point (X, P) is terminal of type cA/n or isolated cDV .

Proof. The only thing we have to prove is the last statement. By the Inversion of Adjunction [Sho93, §3], [Kol92, Ch. 16] the pair (X, H) is PLT. Since H is Cartier and (X, P) is isolated, it is terminal. Clearly, we may assume that (H, P) is not Du Val. Let $F \in |-K_X|$ be a general member. Then $F|_H$ is a general member of $|-K_H|$. Since (H, P) is cyclic quotient (by Proposition 2.2.2), $(H, F|_H)$ is LC. Again by the Inversion of Adjunction the pair $(X, H + F)$ is also LC. But this means that (F, P) is of type A and so (X, P) is of type cA/n . $\square$

2.3. Two-dimensional contractions. The following fact is easy and well-known (see, e.g., [Pro01, Lemma 7.1.11]).

2.3.1. Lemma. Let $v : S \rightarrow R \ni o$ be a rational curve fibration germ over a smooth curve and let $C := v^{-1}(o)_{\text{red}}$. If the pair (S, C) is PLT, then there is an analytic isomorphism

$$S \simeq (\mathbb{P}^1 \times \mathbb{C})/\mu_m(1, a),$$

where $\gcd(a, m) = 1$. The graph $\Delta(S, C)$ is of the form

$$\overset{a_r}{\circ} \text{ --- } \dots \text{ --- } \overset{a_1}{\circ} \text{ --- } \bullet \text{ --- } \overset{b_1}{\circ} \text{ --- } \dots \text{ --- } \overset{b_s}{\circ}$$

where $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate strings.

2.3.2. Lemma ([Sho93, Th. 6.9], [Kol92, Prop. 12.3.1-2]). *Let $v : S \rightarrow R$ be a rational curve fibration germ over a smooth curve and let Δ be an effective $\mathbb{Q}$ -divisor on S such that $K_S + \Delta \equiv 0$ over R . Assume the locus of log canonical singularities $LCS(S, \Delta)$ of (S, Δ) is not connected near a fiber $v^{-1}(o)$, $o \in R$. Then near $v^{-1}(o)$ the pair (S, Δ) is PLT and $\lfloor \Delta \rfloor$ is a disjoint union of two sections.*

2.3.3. Lemma. *Let C be a smooth complete curve contained in a normal surface H . Assume that the pair (H, C) is not PLT at some point, say $P \in C$, and $(K_H + C) \cdot C < 0$. Then*

- (i) *H has at most two singular points on C .*
- (ii) *If H is singular at a point $Q \in C$ and $Q \neq P$, then the pair (H, C) is PLT at Q . The dual graph $\Delta(H, C)$ for the minimal resolution of (H, C) at Q is of the following form:*

$$\bullet \text{---} \underbrace{\circ \text{---} \cdots \text{---} \circ}_{(H, Q)}$$

If moreover (H, Q) is a Gorenstein point, then it is Du Val.

Proof. By the Inversion of Adjunction [Sho93, §3], [Kol92, Ch. 16] one has $(K_H + C)|_C = K_C + \text{Diff}_C(0)$, where $\text{Diff}_C(0)$ is a $\mathbb{Q}$ -divisor with support at $C \cap \text{Sing}(H)$. Moreover, the multiplicity of $\text{Diff}_C(0)$ at every point of $C \cap \text{Sing}(H)$ is at least $1/2$ and its multiplicity at P is at least 1. Since $\deg \text{Diff}_C(0) \leq -\deg K_C = 2$ the assertion of (i) follows. As for (ii), we see that the multiplicity of $\text{Diff}_C(0)$ at Q is less than 1. Again by the Inversion of Adjunction the pair (H, C) is PLT at Q . The rest follows from the classification of surface PLT pairs (see, e.g., [Kol92, Ch. 3]). $\square$

2.4. Lemma. *Let (X, C) be an extremal curve germ and let $f : (X, C) \rightarrow (Z, o)$ be the corresponding contraction. Assume that a member $H \in |\mathcal{O}_X|_C$ is normal. If (X, C) is a $\mathbb{Q}$ -conic bundle germ, then H has only rational singularities.*

Proof. The assertion follows from the observation that $H \rightarrow f(H)$ is a rational curve fibration. $\square$

2.5. Theorem ([Mor88, 7.3], [MP08, 1.3.7]). *Let (X, C) be an extremal curve germ of type (IA) or (IA $^\vee$) and let $P \in X$ be the non-Gorenstein point. Then a general member $F \in |-K_X|$ does not contain C and has only Du Val singularity of type A at P .*

2.6. Proposition. *Let $f : (X, C) \rightarrow (Z, o)$ be a contraction from a threefold with only terminal singularities such that C is a (not necessarily irreducible) curve and $-K_X$ is ample. Let $F \in |-K_X|$ be a general member. Assume that $F \cap C$ is a point P such that (F, P) is a Du Val singularity of type A. Then, for a general member $H \in |\mathcal{O}_X|_C$, the pair $(X, F + H)$ is LC.*

If f is birational, then so is the pair $(Z, F_Z + T)$, where $F_Z = f(F) \in |-K_Z|$ and $T := f(H) \in |\mathcal{O}_Z|$. In this case, (T, o) is a cyclic quotient singularity.

Proof. First we consider the case where f is birational (this case was considered in [Tzi05]). Then $(F_Z, o) \simeq (F, P)$ is a Du Val singularity of type A. Let T be a general hyperplane section of (Z, o) . Then $T \cap F_Z$ is general hyperplane section of (F_Z, o) . Clearly, $T \cap F_Z = \Gamma_1 + \Gamma_2$ for some irreducible curves Γ_i and the pair $(F_Z, \Gamma_1 + \Gamma_2)$ is LC. By the Inversion of Adjunction so is the pair $(Z, F_Z + T)$. Hence

$(T, \Gamma_1 + \Gamma_2)$ is LC and (T, o) is a cyclic quotient singularity (see, e.g., [Kol92, Ch. 3]). Take $H := f^*T$. Then $K_X + F + H = f^*(K_Z + F_Z + T)$, i.e., the contraction f is $K_X + F + H$ -crepant. Hence the pair $(X, F + H)$ is LC.

Now consider the case where Z is a surface. First we claim that $(X, F + H)$ is LC near F . Consider the restriction $\varphi = f_F : (F, P) \rightarrow (Z, o)$. Let $\Xi \subset Z \simeq \mathbb{C}^2$ be the branch divisor of φ . By the Hurwitz formula we can write $K_F = \varphi^*(K_Z + \frac{1}{2}\Xi)$. Hence,

$$K_F + H|_F = \varphi^*\left(K_Z + \frac{1}{2}\Xi + T\right).$$

Using this and the Inversion of Adjunction we get the following equivalences: $(X, F + H)$ is LC near $F \iff (F, H|_F = \varphi^*T)$ is LC $\iff (Z = \mathbb{C}^2, \frac{1}{2}\Xi + T)$ is LC. Thus it is sufficient to show that $(Z, \frac{1}{2}\Xi + T)$ is LC.

Let $\xi(u, v) = 0$ be the equation of $\Xi \subset \mathbb{C}^2$. Then (F, P) is given by the equation $w^2 = \xi(u, v)$ in $\mathbb{C}_{u,v,w}^3$. By the classification of Du Val singularities we can choose coordinates u, v so that

$$\xi = u^2 + v^{n+1}.$$

Take $T := \{v - u = 0\}$. Then $\text{ord}_0 \xi(u, v)|_T = 2$. By the Inversion of Adjunction the pair $(Z, T + \frac{1}{2}\Xi)$ is LC.

Thus we have shown that $(X, F + H)$ is LC near F . Assume that $(X, F + H)$ is not LC at some point $Q \in C$. By the above, $Q \notin F$. Note that H is smooth outside of C by Bertini's theorem.

Assume that H is not normal. Let $\nu : H' \rightarrow H$ be the normalization and let $C' := \nu^{-1}(C)_{\text{red}}$. Write

$$K_{H'} + \text{Diff}_H(F) = \nu^*(K_X + H + F) \sim 0.$$

Here $\text{Diff}_H(F) = C' + \nu^{-1}(F|_H)$, where $C' = \nu^{-1}(C)$. By the Inversion of Adjunction C' is reduced and $(H', C' + \nu^{-1}(F|_H))$ is not LC at $\nu^{-1}(Q)$. Now we can apply Lemma 2.3.2 to $(H', C' + \nu^{-1}(F|_H) - \varepsilon v^*(o))$. $\square$

2.6.1. Corollary. *Under the assumptions of 2.6, if H is not normal, then there is an analytic isomorphism $(H, P) \simeq \{x'_1 x'_2 = 0\} / \mu_m(a, -a, 1)$.*

Proof. Let $\pi : (X^\sharp, P^\sharp) \rightarrow (X, P)$ be the index-one cover and let $H^\sharp := \pi^*H$, $F^\sharp := \pi^*F$. Then the pair $(X^\sharp, H^\sharp + F^\sharp)$ is LC.

Assume that (X, P) is not a cyclic quotient singularity. One can choose a μ_m -equivariant embedding $X^\sharp \subset \mathbb{C}_{x_1, \dots, x_4}^4$ so that $\text{wt}(x_1, \dots, x_4) \equiv (a, -a, 1, 0) \pmod{m}$ and $X^\sharp$ is given by the equation $x_1 x_2 = \phi(x_3^m, x_4)$, where $\text{ord}_0 \phi \geq 2$. For some hypersurfaces $D = \{\xi = 0\}$ and $S = \{\psi = 0\}$ in $\mathbb{C}_{x_1, \dots, x_4}^4$ we have $H^\sharp = D \cap X^\sharp$ and $F^\sharp = S \cap X^\sharp$. By the Inversion of Adjunction the pair $(\mathbb{C}^4, X^\sharp + D + S)$ is LC. On the other hand, by blowing up the origin we get an exceptional divisor of discrepancy

$$a(E, X^\sharp + D + S) = 3 - 2 - \text{ord}_0 \xi - \text{ord}_0 \psi \geq -1.$$

Hence, $\text{ord}_0 \xi = 1$. Since ξ is an μ_m -invariant, it contains the term x_4 . Thus $\xi = x_4 - \xi'$, where $\text{ord}_0 \xi' \geq 2$. Then $H^\sharp$ is given by two equations $x_1 x_2 = \phi(x_3^m, \xi')$ and $x_4 = \xi'$. By changing coordinates we get what we need.

Now assume that (X, P) is a cyclic quotient singularity. Then $X^\sharp \simeq \mathbb{C}^3$. Again one can choose a coordinate system x_1, x_2, x_3 in $\mathbb{C}^3$ so that $\text{wt}(x_1, x_2, x_3) \equiv (a, -a, 1) \pmod{m}$. Let ξ be the equation of $H^\sharp$. By blowing up the origin we get

$\text{ord}_0 \xi \leq 2$. On the other hand, ξ is an invariant. Hence, ξ contains the term $x_1 x_2$ (possibly up to permutations of coordinates if $a \equiv \pm 1$). $\square$

3. DEFORMATIONS OF 3-DIMENSIONAL DIVISORIAL CONTRACTIONS

In this section we recall and set up deformation tools to study extremal curve germs.

3.1. Theorem. *Let $f : (X, C) \rightarrow (Z, o)$ be a 3-dimensional divisorial extremal curve germ, where C is not necessarily irreducible, and let E be its exceptional locus. Then the divisorial part of E is a $\mathbb{Q}$ -Cartier divisor. If furthermore C is irreducible, then E is $\mathbb{Q}$ -Cartier and (Z, o) is a terminal singularity.*

3.2. Theorem (cf. [KM92, (11.4)], [MP08, (6.2)]). *Let $f : (X, C) \rightarrow (Z, o)$ be an extremal divisorial (resp. flipping, $\mathbb{Q}$ -conic bundle) curve germ, where C is not necessarily irreducible. Let $\pi : \mathcal{X} \rightarrow (\mathbb{C}_\lambda^1, 0)$ be a flat deformation of $X = \mathcal{X}_0 := \pi^{-1}(0)$ over a germ $(\mathbb{C}_\lambda^1, 0)$ with a flat closed subspace $\mathcal{C} \subset \mathcal{X}$ such that $C = \mathcal{C}_0$. Then there exist a flat deformation $\mathcal{Z} \rightarrow (\mathbb{C}_\lambda^1, 0)$ and a proper $\mathbb{C}_\lambda^1$ -morphism $\mathfrak{f} : \mathcal{X} \rightarrow \mathcal{Z}$ such that $f = \mathfrak{f}_0$ and $\mathfrak{f}_\lambda : (\mathcal{X}_\lambda, \mathfrak{f}_\lambda^{-1}(o_\lambda)_{\text{red}}) \rightarrow (\mathcal{Z}_\lambda, o_\lambda)$ is a divisorial (resp. flipping, $\mathbb{Q}$ -conic bundle) extremal curve germ for every small λ , where $o_\lambda := \mathfrak{f}_\lambda(\mathcal{C}_\lambda)$.*

3.2.1. Corollary. *Let $f : (X, C) \rightarrow (Z, o)$ be an extremal divisorial curve germ, where C is not necessarily irreducible. Let $P^{(1)}, \dots, P^{(r)} \in X$ be singular points. Let $(X_\lambda, P_\lambda^{(i)}) \supset (C_\lambda, P_\lambda^{(i)})$ be a set of local 1-parameter analytic deformations of $(X, P^{(i)}) \supset (C, P^{(i)})$. Then it extends to a 1-parameter analytic deformation $X_\lambda \supset C_\lambda \supset \{P_\lambda^{(1)}, \dots, P_\lambda^{(r)}\}$ of global $X \supset C \supset \{P^{(1)}, \dots, P^{(r)}\}$ in the sense that there exist a flat deformation $\mathcal{Z} \rightarrow (\mathbb{C}_\lambda^1, 0)$ and a proper $\mathbb{C}_\lambda^1$ -morphism $\mathfrak{f} : \mathcal{X} \rightarrow \mathcal{Z}$ such that $f = \mathfrak{f}_0$ and $\mathfrak{f}_\lambda : (\mathcal{X}_\lambda, \mathfrak{f}_\lambda^{-1}(o_\lambda)_{\text{red}}) \rightarrow (\mathcal{Z}_\lambda, o_\lambda)$ is a divisorial extremal curve germ for every small λ , where $o_\lambda := \mathfrak{f}_\lambda(\mathcal{C}_\lambda)$.*

We need the following easy lemma which can be found in [Bin81, (9.3)] (without proof).

3.3. Lemma. *Let $p : \mathcal{D} \rightarrow \mathcal{X} \supset \ell$ be an arbitrary analytic morphism and let $\ell \subset X$ be a compact subset such that $p^{-1}(\ell)$ is compact. Then there exist open subsets $W \supset p^{-1}(\ell)$ of $\mathcal{D}$ and $V \supset p(W)$ of $\mathcal{X}$ such that $p|_W : W \rightarrow V$ is proper and $p(W)$ is an analytic subset of V .*

Proof. There is an open subset $U \supset p^{-1}(\ell)$ of $\mathcal{D}$ such that $\bar{U}$ is compact (and U is open and closed in $\mathcal{D} \setminus \partial U$). Since $p(\partial U)$ is a closed set disjoint from ℓ , there is an open set $V \supset \ell$ such that $\bar{V}$ is disjoint from $p(\partial U)$. Then $p^{-1}(\bar{V})$ is disjoint from ∂U . Hence $W := U \cap p^{-1}(\bar{V})$ is an open and closed subset of $p^{-1}(\bar{V})$ and is $\bar{W} \subset U$ is compact. Hence $p|_W : W \rightarrow V$ is proper. This means that $p(W)$ is an analytic subset of V . $\square$

The following is the key step in the proof of 3.1 and 3.2.

3.4. Proposition. *Let $f : (X, C) \rightarrow (Z, o)$ be a divisorial extremal curve germ, where C is not necessarily irreducible. Let $\bar{\pi} : \bar{\mathcal{X}} \rightarrow (\mathbb{C}_\lambda^1, 0)$ be a flat deformation of $X = \bar{\mathcal{X}}_0 := \bar{\pi}^{-1}(0)$ over a germ $(\mathbb{C}_\lambda^1, 0)$.*

(i) *Let $\bar{\mathcal{X}}^\wedge$ be the completion of $\bar{\mathcal{X}}$ along $\lambda = 0$. Then $f : X \rightarrow Z$ extends to a contraction $\mathfrak{f}^\wedge : \bar{\mathcal{X}}^\wedge \rightarrow \mathcal{Z}^\wedge$.*

(ii) Let n be an arbitrary positive integer. Then there exist flat deformations $\pi : \mathcal{X} \rightarrow (\mathbb{C}_\lambda^1, 0)$ and $\mathcal{Z} \rightarrow (\mathbb{C}_\lambda^1, 0)$ and a proper $\mathbb{C}_\lambda^1$ -morphism $\mathfrak{f} : \mathcal{X} \rightarrow \mathcal{Z}$ such that $\pi_{(n)} \simeq \bar{\pi}_{(n)}$, $f = \mathfrak{f}_0$, and $\mathfrak{f}_\lambda : \mathcal{X}_\lambda \rightarrow \mathcal{Z}_\lambda$ is a divisorial contraction (which contracts a divisor to a curve) for every small λ , where $\mathcal{A}_{(i)} := \mathcal{A} \times_{\mathbb{C}_\lambda^1} \text{Spec } \mathbb{C}[[\lambda]]/(\lambda^{i+1})$ for any object $\mathcal{A}$ over $\mathbb{C}_\lambda^1$ and $i \geq 0$.

Proof. Let $\phi \in H^0(X, \mathcal{O}_X)$ be a general section vanishing on C and H (resp. H_Z) the member of $|\mathcal{O}_X|$ (resp. $|\mathcal{O}_Z|$) defined by ϕ (resp. $f_*\phi$). We note that H (resp. H_Z) is smooth outside C (resp. o) and f induces an isomorphism $H \setminus C \simeq H_Z \setminus \{o\}$.

Then as in [KM92, (11.3)–(11.4)], the miniversal deformation spaces $\text{Def}(H)$ and $\text{Def}(H_Z)$ exist as analytic spaces and f induces a complex analytic morphism $\text{Def}(f, H) : \text{Def}(H) \rightarrow \text{Def}(H_Z)$. Let $\phi : X \rightarrow \mathbb{C}_s^1$ be the morphism defined by $s = \phi$. This morphism is a flat family of H over $\mathbb{C}_s^1$. Thus we have an induced morphism $\bar{w} : \mathbb{C}_s^1 \rightarrow \text{Def}(H)$, i.e. an element $\bar{w} \in \text{Hom}(\mathbb{C}_s^1, \text{Def}(H))$. Furthermore X , Z , and f can be reconstructed by the morphism $\bar{w} : (\mathbb{C}_s^1, 0) \rightarrow \text{Def}(H)$. Our goal is to construct the following morphism extending $\bar{w}$.

$$w : (\mathbb{C}_{s,\lambda}^2, 0) \longrightarrow \text{Def}(H)$$

Since $R^1 f_* \mathcal{O}_X = 0$, the section ϕ extends to a formal section $\hat{\phi}$ on the completion $\bar{\mathcal{X}}^\wedge$ of $\bar{\mathcal{X}}$ along X . This proves (i). We thus see that $\bar{w} \in \text{Hom}(\mathbb{C}_s^1, \text{Def}(H))$ extends to $\hat{w} \in \text{Hom}((\mathbb{C}_{s,\lambda}^2, 0)^\wedge, \text{Def}(H))$, where $(\mathbb{C}_{s,\lambda}^2, 0)^\wedge$ is the completion of $(\mathbb{C}_{s,\lambda}^2, 0)$ along $\{\lambda = 0\}$. Then by [Art68, Theorem 1.5, (i)], $\hat{w}$ can be approximated by an analytic extension $w \in \text{Hom}((\mathbb{C}_{s,\lambda}^2, 0), \text{Def}(H))$ of $\bar{w}$. This gives us a flat family $\mathcal{X}$ over $\mathbb{C}_\lambda^1$ approximating $\bar{\mathcal{X}}$.

It remains to settle divisoriality. Arbitrarily close to C there is an f -exceptional curve $\ell \simeq \mathbb{P}^1$ such that $N_{\ell/X} \simeq \mathcal{O}_\ell \oplus \mathcal{O}_\ell(-1)$ which sweep out an f -exceptional divisor of X . Hence $N_{\ell/\mathcal{X}} \simeq \mathcal{O}_\ell^{\oplus 2} \oplus \mathcal{O}_\ell(-1)$ and there are no obstructions to deforming these ℓ out to $\mathcal{X}_\lambda$ and hence $\mathfrak{f}_\lambda$ contracts a divisor. This proves the statement (ii) of our proposition. $\square$

Proof of Theorem 3.1. Let $P^{(1)}, \dots, P^{(r)} \in X$ be singular points. As in [Mor88, Appendix 1b] one can see that every local deformation of singularities extends to a deformation of global X . For every terminal singularity $(X, P^{(i)})$ we take a $\mathbb{Q}$ -smoothing, a deformation whose general member has only cyclic quotient singularities [Rei87, (6.4)]. By the above, there exists a one-parameter deformation $\mathcal{X}$ over a disk in $\mathbb{C}_\lambda^1$ such that $\mathcal{X}_0 \simeq X$ and, for small $\lambda \neq 0$, the fiber $\mathcal{X}_\lambda$ has only terminal cyclic quotient singularities. Then we apply Proposition 3.4, (ii). In notation of Proposition 3.4, there exists a divisorial contraction $\mathfrak{f} : \mathcal{X} \rightarrow \mathcal{Z}$ contracting a divisor $\mathcal{E}$ (the divisorial part of the exceptional locus) to a surface on $\mathcal{Z}$ and, for small $\lambda \neq 0$, the fiber $\mathcal{X}_\lambda$ also has only terminal cyclic quotient singularities because at every singular point P of X the local germ of $\mathcal{X}$ at P can be approximated by one of $\mathcal{X}$ to an arbitrarily high order of λ .

Let $P \in X = \mathcal{X}_0$ be a singular point and let $(X^\sharp, P^\sharp)$ be the index-one cover. Then the local deformation $(\mathcal{X}, P)$ is induced by a deformation $(\mathcal{X}^\sharp, P^\sharp)$ of $(X^\sharp, P^\sharp)$ (cf. [Ste88, The last paragraph of §6]). Since the germ $(X^\sharp, P^\sharp)$ is a hypersurface singularity [Rei83], so is $(\mathcal{X}^\sharp, P^\sharp)$. Moreover, the singularity $(\mathcal{X}^\sharp, P^\sharp)$ is isolated. Hence by [Gro68, Exp. XI, Corollary 3.14] the variety $\mathcal{X}^\sharp$ is factorial at $P^\sharp$ and so $\mathcal{X}$ is $\mathbb{Q}$ -factorial at P . In particular, $\mathcal{E}$ is a $\mathbb{Q}$ -Cartier divisor. Thus $\mathcal{E}|_X = E$ on $X \setminus C$. If moreover C is irreducible, then $\rho(X) = 1$ (see [Mor88, (1.3)]) and so

$K_X \sim_{\mathbb{Q}} \mathcal{E}|_X$. Hence, $\mathcal{E}|_X$ is negative on C and $\mathcal{E}|_X \supset C$. This implies that $E = \mathcal{E}|_X$ and it is also $\mathbb{Q}$ -Cartier. $\square$

Proof of Theorem 3.2. The flipping case follows from [KM92, (11.4)] and the $\mathbb{Q}$ -conic bundle case from [MP08, (6.2)]. So, we assume that f is divisorial. Let $E \subset X$ be the exceptional divisor of f and let E_i 's be its irreducible components. Then, for each i , $B_i := f(E_i) \subset Z$ is an irreducible curve passing through o .

First, we treat the case where C is irreducible. Then by Theorem 3.1 E is a $\mathbb{Q}$ -Cartier divisor and $Z \ni o$ is a terminal singularity.

For each E_i , choose a smooth fiber ℓ'_i of $E_i \rightarrow B_i$ and let $[\ell'_i]$ degenerate to $[\ell_i]$ lying over o in the Douady space of X/Z . We assume each $[\ell'_i]$ is chosen arbitrarily close to $[\ell_i]$. Consider the closed subspace A' of the Douady space of X/Z parametrizing all compact subspaces $F \subset X$ with $\text{Supp } F \subset C$. Then each irreducible component of A' is compact, cf. [Fuj79], and let A be the smallest open and closed subset of A' containing all $[\ell_i]$. Thus A is also compact. Then we work on a sufficiently small neighborhood $\mathcal{D}'$ of A in the Douady space of $\mathcal{X}/\mathbb{C}_\lambda^1$ such that $\mathcal{D}' \ni [\ell'_i]$ for each i .

We note that $\mathcal{X}$ is smooth along each ℓ'_i and $N_{\ell'_i/\mathcal{X}} \simeq \mathcal{O}_{\ell'_i}^{\oplus 2} \oplus \mathcal{O}_{\ell'_i}(-1)$. Hence $\mathcal{D}'$ is smooth of dimension 2 at each $[\ell'_i]$. Let $\mathcal{D} \subset \mathcal{D}'$ be the smallest one among the union of the irreducible components of $\mathcal{D}'$ such that $\mathcal{D} \ni [\ell'_i]$ for all i . Then $\mathcal{D}$ is a two-dimensional closed subspace of $\mathcal{D}'$.

Let $\mathcal{T} \subset \mathcal{X} \times_{\mathbb{C}_\lambda^1} \mathcal{D}$ be the universal closed subspace parametrized by $\mathcal{D}$ with two projections $\pi : \mathcal{T} \rightarrow \mathcal{D}$ and $p : \mathcal{T} \rightarrow \mathcal{X}$.

We note that $p^{-1}(C) \subset A$ and it is compact because the variety $\mathcal{X}_0 = X$ has a divisorial contraction to Z , C is the fiber over $o \in Z$, and $\pi^{-1}(t)$ does not intersect C for $t \notin A$.

Let $\mathcal{E} := p(\mathcal{T}) \subset \mathcal{X}$ be the image of the proper morphism p and it is an analytic subset by Lemma 3.3. We also denote by $p : \mathcal{T} \rightarrow \mathcal{E}$ the morphism induced by p and let $p : \mathcal{T} \xrightarrow{p'} \bar{\mathcal{T}} \xrightarrow{\bar{p}} \mathcal{E}$ be the Stein factorization of p so that $p'_* \mathcal{O}_{\mathcal{T}} = \mathcal{O}_{\bar{\mathcal{T}}}$.

3.4.1. Claim. $\mathcal{E}$ is a $\mathbb{Q}$ -Cartier divisor.

Proof. Let $\mathcal{X}^\wedge$ be the completion of $\mathcal{X}$ along $\lambda = 0$. By Proposition 3.4, (i) the morphism $f : X \rightarrow Z$ extends to a contraction $\mathfrak{f}^\wedge : \mathcal{X}^\wedge \rightarrow \mathcal{Z}^\wedge$, where $\mathcal{Z}^\wedge$ is $\mathbb{Q}$ -Gorenstein [Ste88, Corollary 10] because Z is terminal. Comparing $K_{\mathcal{X}^\wedge}$ and $\mathfrak{f}^{\wedge*} K_{\mathcal{Z}^\wedge}$ we see that there is an effective $\mathbb{Q}$ -Cartier divisor $\mathcal{F}^\wedge \sim_{\mathbb{Q}} K_{\mathcal{X}^\wedge} - \mathfrak{f}^{\wedge*} K_{\mathcal{Z}^\wedge}$ on $\mathcal{X}^\wedge$ such that $\mathcal{F}^\wedge|_{\mathcal{X}^\wedge} = E^\wedge$ and $\mathcal{F}^\wedge = \mathcal{E}^\wedge$ outside of $C^\wedge$. Hence $\mathcal{F}^\wedge = \mathcal{E}^\wedge$. $\square$

Now we define a morphism $q : \mathcal{D} \rightarrow \mathcal{B}$ such that $q(p^{-1}(C))$ is one point as follows. Take a general point ζ of C and take a small 3-dimensional disk $(\Delta^3, 0)$ centered at ζ and transversal to C at ζ . Then the Cartier divisor Δ^3 in a neighborhood of C induces a Cartier divisor of $\mathcal{T}$ finite and flat over $\mathcal{D}$. Let d be the degree of $p^{-1}(\Delta^3)/\mathcal{D}$. Then $x \in \mathcal{D} \mapsto \pi^{-1}(x) \cap p^{-1}(\Delta^3)$ associates to x a 0-cycle of degree d on Δ^3 and we have thus a required morphism $q : \mathcal{D} \rightarrow \mathcal{B} := S^d(\Delta^3)$ such that $q(p^{-1}(C))$ is the 0-cycle $d \cdot [0]$.

We claim that we have a proper morphism $r : \bar{\mathcal{T}} \rightarrow \mathcal{B}$ making the following diagram commutative

$$\begin{array}{ccccc} \mathcal{T} & \xrightarrow{p'} & \bar{\mathcal{T}} & \xrightarrow{\bar{p}} & \mathcal{E} \\ \downarrow \pi & & \downarrow r & & \\ \mathcal{D} & \xrightarrow{q} & \mathcal{B} & & \end{array}$$

Indeed since $q(\pi(p^{-1}(C)))$ is one-point $d \cdot [0]$, we can shrink $\mathcal{E}$ so that $q(\mathcal{D})$ is contained in a Stein open neighborhood of $d \cdot [0]$. Hence the morphism $\mathcal{T} \rightarrow \mathcal{B}$ factors through $p' : \mathcal{T} \rightarrow \bar{\mathcal{T}}$ and the claim is proved.

We claim that $p, p', \bar{p}$ are isomorphisms over every ℓ'_i and in particular $\bar{p}$ is finite and bimeromorphic. Indeed by $N_{\ell'_i/\mathcal{X}} \simeq \mathcal{O}_{\ell'_i}^{\oplus 2} \oplus \mathcal{O}_{\ell'_i}(-1)$, p is an isomorphism near $\pi^{-1}([\ell'_i])$ and by the divisorial contraction on $X = \{\lambda = 0\} \subset \mathcal{X}$ one has $p^{-1}(\ell'_i) = \pi^{-1}([\ell'_i])$. These settle the claim.

Let $\mathfrak{c} := \mathcal{H}om_{\mathcal{O}_{\bar{\mathcal{T}}}}(\mathcal{O}_{\bar{\mathcal{T}}}, \mathcal{O}_{\mathcal{T}})$ be the conductor of $\bar{p}$ and let $V(\mathfrak{c}) \subset \bar{\mathcal{T}}$ be the locus defined by $\mathfrak{c}$. Then we claim that $r(V(\mathfrak{c}))$ is finite over $\mathbb{C}_{\lambda}^1$. Indeed this is obvious since $r(V(\mathfrak{c})) \not\supset q([\ell'_i])$ and the fiber of $r(V(\mathfrak{c}))$ over $\{\lambda = 0\}$ is a finite set.

Let $J \subset \mathcal{O}_{\mathcal{B}}$ be an arbitrary sheaf of ideals such that $J\mathcal{O}_{\bar{\mathcal{T}}} \subset \mathfrak{c}$ and $V(J)$ is finite over $\mathbb{C}_{\lambda}^1$. By [Bin81, Theorem (6.1)] we have the following diagram

$$\begin{array}{ccc} V(J) & \longrightarrow & \mathbb{C}_{\lambda}^1 \\ \downarrow & & \downarrow \\ \mathcal{B} & \xrightarrow{q'} & \mathcal{E}' \end{array}$$

where $\mathcal{E}' := \mathcal{B} \amalg_{V(J)} \mathbb{C}_{\lambda}^1$ is the amalgamated sum (coproduct) of $\mathcal{B}$ and $\mathbb{C}_{\lambda}^1$ over $V(J)$ and $q' : \mathcal{B} \rightarrow \mathcal{E}'$ is a bimeromorphic finite morphism. Since $\mathfrak{c}$ is the conductor of $\bar{p}$, we have

$$\mathcal{E} = \bar{\mathcal{T}} \amalg_{V_{\bar{\mathcal{T}}}(\mathfrak{c})} V_{\mathcal{E}}(\mathfrak{c})$$

and the following commutative diagram.

$$\begin{array}{ccc} V_{\bar{\mathcal{T}}}(\mathfrak{c}) & \longrightarrow & V_{\mathcal{E}}(\mathfrak{c}) \\ \downarrow & & \downarrow \\ \bar{\mathcal{T}} & \xrightarrow{\bar{p}} & \mathcal{E} \end{array}$$

These two diagrams fit into a big one which allows us to define an induced morphism $\eta : \mathcal{E} \rightarrow \mathcal{E}'$:

$$\begin{array}{ccccccc} V_{\bar{\mathcal{T}}}(\mathfrak{c}) & \longrightarrow & V_{\mathcal{E}}(\mathfrak{c}) & & & & \\ \downarrow & \searrow & \downarrow & \searrow & r & & \\ \bar{\mathcal{T}} & \longrightarrow & \mathcal{E} & \xrightarrow{\eta} & V(J) & \longrightarrow & \mathbb{C}_{\lambda}^1 \\ & \searrow & & \searrow & \downarrow & & \downarrow \\ & & \mathcal{B} & \xrightarrow{q'} & \mathcal{E}' & & \end{array}$$

Finally we have the following commutative diagram:

$$\begin{array}{ccccc} \mathcal{T} & \xrightarrow{p'} & \bar{\mathcal{T}} & \xrightarrow{\bar{p}} & \mathcal{E} \\ \downarrow \pi & & \downarrow r & & \downarrow \eta \\ \mathcal{D} & \xrightarrow{q} & \mathcal{B} & \xrightarrow{q'} & \mathcal{E}' \end{array}$$

For any $i, j \geq 0$ the sheaf $\mathcal{O}_{i\mathcal{E}}(-j\mathcal{E})$ denotes the quotient $\mathcal{O}_{\mathcal{X}}(-j\mathcal{E})/\mathcal{O}_{\mathcal{X}}(-(i+j)\mathcal{E})$.

3.4.2. Claim. *For any $i, j \geq 0$ we have $R^1\eta_*\mathcal{O}_{i\mathcal{E}}(-j\mathcal{E}) = 0$. Therefore, the following sequence*

$$0 \longrightarrow \eta_*\mathcal{O}_{i\mathcal{E}}(-j\mathcal{E}) \longrightarrow \eta_*\mathcal{O}_{(i+j)\mathcal{E}} \longrightarrow \eta_*\mathcal{O}_{j\mathcal{E}} \longrightarrow 0.$$

is exact.

Proof. By the Kawamata-Viehweg vanishing [Nak87, Theorem 3.6] $R^1f_*\mathcal{O}_X(-kE) = 0$ for $k \geq 0$. Then from the exact sequence

$$0 \longrightarrow \mathcal{O}_X(-(i+j)E) \longrightarrow \mathcal{O}_X(-jE) \longrightarrow \mathcal{O}_{iE}(-jE) \longrightarrow 0$$

we see that $R^1f_*\mathcal{O}_{iE}(-jE) = 0$ for $i, j \geq 0$. Now we assert that the following sequence

$$0 \longrightarrow \mathcal{O}_{i\mathcal{E}}(-j\mathcal{E}) \xrightarrow{\lambda} \mathcal{O}_{i\mathcal{E}}(-j\mathcal{E}) \longrightarrow \mathcal{O}_{iE}(-jE) \longrightarrow 0.$$

is exact for $i, j \geq 0$. Recall that the space $\mathcal{X}$ is $\mathbb{Q}$ -Gorenstein [Ste88, The last paragraph of §6]. Consider the index-one cover $\nu : (\mathcal{X}^\sharp, P^\sharp) \rightarrow (\mathcal{X}, P)$ with respect to $\mathcal{E}$ at an arbitrary point $P \in X$. Since the map ν is étale in codimension two, both $\mathcal{X}^\sharp$ and $X^\sharp := \nu^{-1}(X)$ are terminal. The induced divisors $\mathcal{E}^\sharp$ and $E^\sharp$ are Cartier on $\mathcal{X}^\sharp$ and $X^\sharp$, respectively, and $E^\sharp = \mathcal{E}^\sharp|_{X^\sharp}$. Hence the assertion on exactness can be readily checked on $\mathcal{X}^\sharp$. Then by Nakayama's lemma we obtain $R^1\eta_*\mathcal{O}_{i\mathcal{E}}(-j\mathcal{E}) = 0$. $\square$

Fix a positive integer m such that both mE and $m\mathcal{E}$ are Cartier and define a ringed space $\mathcal{E}''$ as a topological space $\mathrm{Spec}_{\mathcal{E}'} \eta_*\mathcal{O}_{\mathcal{E}}$ with the sheaf of rings $\eta_*\mathcal{O}_{m\mathcal{E}}$. Then $\mathcal{E}''$ is a complex space by Claim 3.4.2 and [Bin81, §10].

Now we would like to show that $\mathcal{X}$ has a modification, and in order to do that we would like to check the conditions (1) and (2) of Corollary (8.2) of Bingener [Bin81] for the morphism $\mathcal{X} \supset m\mathcal{E} \rightarrow \mathcal{E}''$ induced by η . The condition (1) is obvious because $-\mathcal{E}$ is ample, and the condition (2) follows from the exact sequence in Claim 3.4.2 with $j = 1$. Thus the desired contraction $\mathfrak{f} : \mathcal{X} \rightarrow \mathcal{Z}$ exists by [Bin81, Corollary (8.2)]. So the proof of the case of irreducible C is completed.

Now we consider the general case, i.e. we assume that C is reducible. Run an analytic MMP on $\mathcal{X}$ in the following way. Every irreducible K -negative curve on the central fiber of X/Z generates an extremal ray on X . By [KM92, (11.7)] flips on X extend to ones on $\mathcal{X}$. So do divisorial contractions by our previous arguments. By Theorem 3.1 we stay in the terminal category. At the end we get $X' \subset \mathcal{X}'/\mathbb{C}_\lambda^1$ such that X' is a minimal model over Z . Moreover, all fibers of $f' : X' \rightarrow Z$ are of dimension ≤ 1 and $-K_{X'}$ is ample over Z outside of the central fiber. Hence, $f' : X' \rightarrow Z$ is a small contraction. Note that $R^1f'_*\mathcal{O}_{X'} = R^1f_*\mathcal{O}_X = 0$. By [KM92, (11.4)] the contraction $f' : X' \rightarrow Z$ extends to $\mathfrak{f}' : \mathcal{X}' \rightarrow \mathcal{Z}$. Thus we have a bimeromorphic map $\mathfrak{f} : \mathcal{X} \dashrightarrow \mathcal{Z}$. By Zariski's Main Theorem this map is actually a proper morphism. This proves Theorem 3.2. $\square$

4. CASE cD/3

In this section we prove Theorem 1.4.

4.1. Setup. Let (X, C) be an extremal curve germ and let $f : (X, C) \rightarrow (Z, o)$ be the corresponding contraction. In particular, f can be flipping. Throughout this section we assume that (X, C) is of type (IA) and the only non-Gorenstein point $P \in (X, C)$ is of type cD/3 [Mor85], [Rei87]. Our arguments here are very similar to those in [KM92, §6]. Note that by Corollary 2.2.4 the point (H, P) is not log terminal for any divisor $H \in |\mathcal{O}_X|_C$. Let $\sigma = (\sigma_1, \dots, \sigma_n)$ be a weight. Below, for a formal power series α in n variables, $\alpha_{\sigma=m}$ means the sum of the monomials in α whose σ -weight is m . Put $\sigma := (1, 1, 2, 3)$. As in [KM92, 6.5], up to coordinate change the point (X, P) is given by

$$\{\alpha(y_1, y_2, y_3, y_4) = 0\} \subset \mathbb{C}_{y_1, y_2, y_3, y_4}^4 / \mu_3(1, 1, 2, 0),$$

where

$$\alpha = y_4^2 + y_3^3 + \delta_3(y_1, y_2) + (\text{terms of degree} \geq 4),$$

$\delta_3(y_1, y_2) = \alpha_{\sigma=3}(y_1, y_2, 0, 0) \neq 0$, $\text{wt } \alpha \equiv 0 \pmod{3}$, and $C^\sharp$ is the y_1 -axis. If $\delta_3(y_1, y_2)$ is squarefree (resp. has a double factor, is a cube of a linear form), then (X, P) is said to be a *simple* (resp. *double*, *triple*) cD/3 point. The general member $F \in |-K_X|$ modulo a coordinate change is given by the equation $y_1 = 0$ (see [Rei87]).

4.2. Lemma. *In the above coordinate system there exists a member $H \in |\mathcal{O}_X|_C$ given by the equation $y_4 = \xi$, where $\xi = \xi(y_1, y_2, y_3)$ is an invariant in the ideal $(y_2, y_3)^3 + y_1(y_2, y_3)$.*

Proof. We have the following exact sequence

$$0 \longrightarrow \omega_X \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_F \longrightarrow 0.$$

If f is a birational contraction, then $R^1 f_* \omega_X = 0$ by the Grauert-Riemenschneider vanishing theorem. Hence any section $\bar{s} \in \mathcal{O}_F$ lifts to a section $s \in f_* \mathcal{O}_X$. So, the assertion is clear in this case. Assume that f is a $\mathbb{Q}$ -conic bundle. Obviously, $\tau := f|_F$ is a double cover. Since $R^1 f_* \omega_X = \omega_Z$ (see [MP08, Lemma 4.1]) and $\omega_F \simeq \mathcal{O}_F$, we have

$$f_* \mathcal{O}_X \longrightarrow \tau_* \omega_F \longrightarrow \omega_Z \longrightarrow 0.$$

The last map is nothing but the trace map $\text{Tr}_{F/Z} : \tau_* \omega_F \rightarrow \omega_Z$. According to [MP09, 2.1-2.2] the induced map

$$f_* \mathcal{O}_X \longrightarrow \tau_* \omega_F / \tau^* \omega_Z$$

is surjective. We may assume that the equation of F in $\mathbb{C}_{y_2, y_3, y_4}^3$ is as follows

$$\beta(y_2, y_3, y_4) := \alpha(0, y_2, y_3, y_4) = y_4^2 + y_3^3 + \delta_3(0, y_2) + (\text{terms of degree} \geq 4).$$

Locally, near $P^\sharp$, the sheaf $\omega_{F^\sharp}$ is generated by

$$\eta := \text{Res} \frac{dy_2 \wedge dy_3 \wedge dy_4}{\beta} = -\frac{dy_2 \wedge dy_3}{\partial \beta / \partial y_4} = \frac{dy_2 \wedge dy_4}{\partial \beta / \partial y_3} = -\frac{dy_3 \wedge dy_4}{\partial \beta / \partial y_2}.$$

Since η is an invariant, it is also a generator of ω_F near P . Further, since Z is smooth, one has

$$\tau^* \Omega_Z^2 = \tau^* \omega_Z \subset \Omega_F^2 \longrightarrow \omega_F.$$

The generators of $\mathcal{O}_{F,P}$ are y_4 , $w := y_2 y_3$, $u := y_2^3$, and $v := y_3^3$ with relations $uv = w^3$ and $y_4^2 + v + u + \dots = 0$. Eliminating v we get three generators y_4 , w , u and one relation $u(u + y_4^2 + \dots) + w^3 = 0$. Hence Ω_F^2 is generated by the elements

$$\begin{aligned} dw \wedge du &= d(y_2 y_3) \wedge d(y_2^3) = 3y_2^3 dy_3 \wedge dy_2, \\ du \wedge dy_4 &= d(y_2^3) \wedge dy_4 = 3y_2^2 dy_2 \wedge dy_4, \\ dw \wedge dy_4 &= d(y_2 y_3) \wedge dy_4 = y_2 dy_3 \wedge dy_4 + y_3 dy_2 \wedge dy_4. \end{aligned}$$

Then Ω_F^2 is contained in ηI , where

$$I := \langle y_3^3 \partial \beta / \partial y_4, y_2^2 \partial \beta / \partial y_3, y_2 \partial \beta / \partial y_2, y_3 \partial \beta / \partial y_3 \rangle \subset (y_2, y_3, y_4)^3.$$

So, $\tau^* \omega_Z \subset (\tau_* \omega_F) I$. Therefore, for some $\xi \in I$ the section $\bar{s} = y_4 - \xi \in \mathcal{O}_F$ lifts to a section $s \in f_* \mathcal{O}_X$. Since

$$s \equiv y_4 \pmod{(y_2, y_3, y_4)^3 + y_1(y_2, y_3, y_4)},$$

one can apply Weierstrass' preparation theorem to get Lemma 4.2. $\square$

4.3. Corollary. *If y_4 is a part of an ℓ -free ℓ -basis of $\text{gr}_C^1 \mathcal{O}$, then a general member $H \in |\mathcal{O}_X|_C$ is normal.*

4.4. Recall that $\ell(P) := \text{len}_{P^\sharp} I^{\sharp(2)} / I^{\sharp 2}$ [Mor88, 9.4.7]. According to [Mor88, 2.16] we have $i_P(1) = \lfloor \ell(P)/3 \rfloor + 1$ and the coordinate system (y_i) can be chosen so that $\alpha \equiv y_1^{\ell(P)} y_i \pmod{(y_2, y_3, y_4)^2}$, where $i \in \{2, 3, 4\}$ and $\ell(P) + \text{wt } y_i \equiv 0 \pmod 3$. Since (X, P) is of type cD/3, we have $\ell(P) > 1$.


Now we are going to prove Theorem 1.4 by considering cases according to the value of $\ell(P)$. We start with the case $\ell(P) = 2$.

4.5. Theorem. *Let the notation and assumptions be as in 4.1. Assume that $\ell(P) = 2$ or, equivalently, $i_P(1) = 1$. Then the following assertions hold.*


4.5.1. *The contraction f is birational; the general member $H \in |\mathcal{O}_X|_C$ and its image $T = f(H) \in |\mathcal{O}_Z|$ are normal and have only rational singularities.*

4.5.2. *If f is flipping (resp. divisorial), then P is not a triple cD/3 point and the dual graph of (H, C) is given as follows with $a = 0$ (resp. $a = 1$).*

(4.5.2.1) *Case of simple cD/3 point P :*



(4.5.2.2) *Case of double cD/3 point P :*




4.5.3. $\text{gr}_C^1 \mathcal{O} = (a) \tilde{\oplus} (-a + P^\sharp)$


We now start the proof of Theorem 4.5.

Proof. Additionally to 4.1 we assume that $\ell(P) = 2$. Then by [Mor88, 2.16] $i_P(1) = 1$ and (in some coordinate system) α satisfies $\alpha \equiv y_1^2 y_2 \pmod{(y_2, y_3, y_4)^2}$. Here $C^\sharp$ is the y_1 -axis as above. Hence, y_3, y_4 form an ℓ -basis of $\text{gr}_C^1 \mathcal{O}$. By Corollary 4.3 H is normal and by Lemma 2.3.3 $H \setminus \{P\}$ can have at most one singular point R which is Du Val. Therefore, X can have at most one type (III) point.

4.5.4. Subcase $\alpha_{\sigma=3}(y_1, y_2, 0, 0)$ is squarefree. By [KM92, 6.7.1] and Lemma 2.3.3 the graph $\Delta(H, C)$ is of the form




We have $a \leq 1$, since the corresponding matrix is negative semi-definite. But then this matrix is negative definite. Hence the contraction f is birational. If $a = 1$, then H is contracted to a singularity $T = f(H)$ of type A_2 . Since T is Gorenstein, f is a divisorial contraction as in 1.4.1. If $a = 0$, that is, P is the only singular point of H , then H is contracted to a singularity $T = f(H)$ with the following dual graph



Let $s \in H^0(X, \mathcal{O}_X)$ be the section defining H . Then $s\mathcal{O}_C \subset \text{gr}_C^1 \mathcal{O}$ is a subbundle outside P since $H \setminus \{P\}$ is smooth. At $P^\sharp$, $s\mathcal{O}_C^\sharp$ is a subbundle of $\text{gr}_C^1 \mathcal{O}^\sharp$ by Lemma 4.2, whence $s\mathcal{O}_C \simeq (0)$ with ℓ -structure. Since $\deg \text{gr}_C^1 \mathcal{O} = 0$ by $i_P(1) = 0$, we have $\text{gr}_C^1 \mathcal{O} = (0) \oplus (P^\sharp)$. Thus f is flipping by [KM92, (6.2.4)].

By 4.5.4 it remains to consider the case where $\alpha_{\sigma=3}(y_1, y_2, 0, 0)$ has a double factor. Note that y_2 divides $\alpha_{\sigma=3}(y_1, y_2, 0, 0)$ because $C^\sharp = (y_1\text{-axis}) \subset X^\sharp$. Since $\ell(P) = 2$, $y_2 y_1^2 \in \alpha$. Then making a coordinate change $y_1 \mapsto y_1 + cy_2$ we get $\alpha_{\sigma=3}(y_1, y_2, 0, 0) = y_1^2 y_2$ and $C^\sharp$ unchanged.

4.5.5. Subcase $\alpha_{\sigma=3}(y_1, y_2, 0, 0) = y_1^2 y_2$ and $\alpha_{\sigma=6}(0, y_2, y_3, 0)$ is squarefree. As above by [KM92, 6.7.2] and Lemma 2.3.3 the graph $\Delta(H, C)$ is of the form



with $a \leq 1$. Again if $a = 1$, then T is Du Val of type D_4 , so f is a divisorial contraction as in 1.4.2. If $a = 0$, then similarly to 4.5.4 the contraction f is flipping (cf [KM92, (6.2.3.2)]). Since $s\mathcal{O}_C^\sharp$ is a subbundle of $\text{gr}_{C^\sharp}^1 \mathcal{O}_C^\sharp$ at $P^\sharp$, as we saw above, it is easy to see 4.5.3.

4.5.6. Subcase $\alpha_{\sigma=3}(y_1, y_2, 0, 0) = y_1^2 y_2$, and $\alpha_{\sigma=6}(0, y_2, y_3, 0)$ has a multiple factor. We will show that this case does not occur. Assume that f is birational. Then the map $H^0(\mathcal{O}_X) \rightarrow H^0(\mathcal{O}_F)$ is surjective [Mor88, Th. 1.2]. Therefore, for any $\lambda \in \mathbb{C}^*$ there is a semi-invariant δ with $\text{wt } \delta = 2$ such that the section $y_4 + \lambda y_2^3 + \delta y_1$ extends to some element $H' \in |\mathcal{O}_X|_C$. After the coordinate change $y'_4 = y_4 + \lambda y_2^3 + \delta y_1$ we see that H' is given by $y'_4 = 0$ and $\alpha' = \alpha(y_1, y_2, y_3, y'_4 - \lambda y_2^3 - \delta y_1)$. Note that $y_4^2 \in \alpha$, $y_4 \notin \alpha$, and α may contain $y_2^3 y_4$. Thus $\alpha'_{\sigma=3}(y_1, y_2, y_3, 0) =$


$\alpha_{\sigma=3}(y_1, y_2, y_3, 0)$ and $\alpha'_{\sigma=6}(0, y_2, y_3, 0) = \alpha_{\sigma=6}(0, y_2, y_3, 0) + (\lambda^2 + c\lambda)y_2^6$ for some $c \in \mathbb{C}$. Hence we may assume that $\alpha_{\sigma=6}(0, y_2, y_3, 0)$ is square free. This contradicts our assumption. (In fact, the above arguments show that the chosen H is not general).

Therefore, f is a $\mathbb{Q}$ -conic bundle. By Lemma 2.4 (H, P) is a rational singularity and by Lemma 4.2 this singularity is analytically isomorphic to

$$\{\gamma(y_1, y_2, y_3) = 0\}/\mu_3(1, 1, 2),$$

where $\gamma(y_1, y_2, y_3) := \alpha(y_1, y_2, y_3, \xi)$, and $C \subset H$ is the image of y_1 -axis. Note that the pair (H, C) is not plt at P . Indeed, otherwise the singularity $\{\gamma = 0\}$ is log terminal (see [Kol92, Cor. 20.4]). Hence it is Du Val. On the other hand, $\text{ord } \gamma > 2$, a contradiction. Let $\sigma' := (1, 1, 2)$. Note that $\gamma_{\sigma'=6}(0, 0, 1) \neq 0$ because $y_3^3 \in \alpha$. Consider the weighted σ' -blowup $\varsigma : \overline{H} \subset \mathbb{C}^3/\mu_3 \rightarrow H \subset \mathbb{C}^3/\mu_3$. Let $\Xi := \varsigma^{-1}(0)_{\text{red}}$. The exceptional divisor $\Theta \subset \overline{H}$ is given in $\Xi \simeq \mathbb{P}(1, 1, 2)$ by the equation $\gamma_{\sigma'=3}(y_1, y_2, 0) = y_1^2 y_2 = 0$. Hence, $\Theta = 2\Theta_1 + \Theta_2$, where Θ_i are irreducible toric divisors in $\mathbb{P}(1, 1, 2)$. The proper transform $\overline{C}$ of C meets $\Xi \simeq \mathbb{P}(1, 1, 2)$ at the point $\{y_2 = y_3 = 0\}$. So, $\overline{C} \cap \Theta_1 = \emptyset$. Since Θ_2 is a smooth reduced component of the Cartier divisor $\Theta = \Xi \cap \overline{H}$ on $\overline{H}$, we see that $\overline{H}$ is smooth at points on $\Theta_2 \setminus \Theta_1$.

In the chart $U_3 \simeq \mathbb{C}^3/\mu_2(1, 1, 1)$ over $\{y_3 \neq 0\}$ we have a new coordinate system $y_1 \mapsto y_1 y_3^{1/3}$, $y_2 \mapsto y_2 y_3^{1/3}$, $y_3 \mapsto y_3^{2/3}$. Here the surface $\overline{H}$ is given by the equation $y_1^2 y_2 + \gamma_{\sigma'=6}(y_1, y_2, 1)y_3 + (\dots)y_3^2 = 0$, where $\gamma_{\sigma'=6}(0, 0, 1) \neq 0$. The origin $O_3 \in \overline{H} \cap U_3$ is a Du Val point of type A_1 . Components Θ_1 and Θ_2 of the exceptional divisor meet each other at O_3 and the pair $(\overline{H}, \Theta_1 + \Theta_2)$ is LC at O_3 . Outside of O_3 , $\overline{H}$ is a hypersurface and has only rational singularities. Therefore, the singularities of $\overline{H}$ are Du Val. Thus the configuration of curves $\overline{C}$, Θ_1 , and Θ_2 on $\overline{H}$ looks as follows:



where $Q_1, \dots, Q_l$ are some Du Val points. By Lemma 2.3.3 the dual graph $\Delta(H, C)$ is of the form

$$(4.5.6.1) \quad \overbrace{\circ \cdots \circ}^a - \underset{C}{\bullet} - \underset{\Theta_2}{\underset{\circ}{\circ}} - \circ - \underset{\Theta_1}{\underset{\circ}{\circ}} - \vdots - \boxed{}$$


where the box on the right hand side indicates some Du Val graphs and the number of these Du Val tails is not important. This configuration forms a fiber of a rational curve fibration. Contracting black vertices successively we obtain

$$(4.5.6.2) \quad \underset{\Theta_2}{\underset{\circ}{\circ}}^{b_2-a-1} - \circ - \underset{\Theta_1}{\underset{\circ}{\circ}}^{b_1} - \vdots - \boxed{}$$


This is again a dual graph of a fiber of a rational curve fibration. Hence $b_2 - a - 1 = 1$ and we further obtain

$$b_1 - 1 \circ - \vdots - \boxed{}$$

Hence $b_1 = 2$ (because the last graph must contain a (-1) -vertex) and so the graph (4.5.6.2) consists of (-2) and (-1) -curves. Furthermore the graph (4.5.6.2) is not a linear chain because the pair (H, C) is not PLT at P . In this situation there is only one possibility (see, e.g., [Pro01, Lemma 7.1.3, 7.1.12]):



Therefore, the original graph (4.5.6.1) is of the form



But then H has only log terminal singularities (see, e.g., [Kol92, Ch. 3]). Hence H has only T-singularities (see 2.2.1) while the right hand side singularity is not of type T (see Proposition 2.2.2), a contradiction. Thus the case 4.5.6 does not occur.

Now the assertion of Theorem 4.5 follows from 4.5.4, 4.5.5, and 4.5.6. This completes our treatment of the case $\ell(P) = 2$. $\square$

4.6. Corollary. *In the notation of 4.1 X has at most one type (III) point.*

Proof. If X has two type (III) points R_1 and R_2 , then by [Mor88, (2.3.3)] and [MP08, (3.1.5)] we have $i_P(1) = i_{R_1}(1) = i_{R_2}(1) = 1$. Then by [Mor88, 2.16] $\ell(P) = 2$. This contradicts Theorem 4.5. $\square$

4.7. Lemma (cf. [KM92, Lemma 6.12]). *If, in the notation of 4.1, X has a type (III) point, then $\ell(P) \leq 4$ and $i_P(1) \leq 2$.*

Proof. Assume that $\ell(P) \geq 5$. As in 4.4 take a coordinate system so that $\alpha \equiv y_1^{\ell(P)} y_i \pmod{(y_2, y_3, y_4)^2}$, where $i \in \{2, 3, 4\}$ and $\ell(P) + \text{wt } y_i \equiv 0 \pmod{3}$. Similarly to the proof of [KM92, Lemma 6.12] we use the deformation $\alpha_\lambda = \alpha + \lambda y_1^{\ell(P)-3} y_i$ (see Theorem 3.2) and get a germ (X_λ, C_λ) with two type (III) points and a point of type cD/3. This contradicts Corollary 4.6. $\square$

For the case $\ell(P) \geq 3$, we are going to prove the following, which settles Theorem 1.4.

4.8. Theorem. *Let the notation and assumptions be as in 4.1. Assume $\ell(P) \geq 3$ or, equivalently, $i_P(1) \geq 2$. Then the following assertions hold.*

4.8.1. $\ell(P) = 3$ or 4 (i.e. $i_P(1) = 2$), and f is birational.

4.8.2. P is a double (resp. triple) cD/3 point if (X, C) is isolated (resp. divisorial).


4.8.3. X is smooth outside of P and there is an ℓ -isomorphism

$$(4.8.3.1) \quad \text{gr}_C^1 \mathcal{O} = ((4 - \ell(P))P^\sharp) \tilde{\oplus} (-1 + 2P^\sharp).$$


4.8.4. For general members $D \in |K_X|$ and $D' \in |K_X|$ (resp. $D' \in |\mathcal{O}_X|_C$), $D \cap D'$ is equal to $4C$ (resp. $3C$) as a 1-cycle.

4.8.5. *The general member $H \in |\mathcal{O}_X|_C$ and its image $T = f(H) \in |\mathcal{O}_Z|$ are normal and have only rational singularities. The dual configuration of (H, C) is as follows.*

(4.8.5.1) *Case of isolated (X, C) :*



(4.8.5.2) *Case of divisorial (X, C) :*



4.8.6. *Conversely if (X, C) is an arbitrary germ of a threefold along $C \simeq \mathbb{P}^1$ with a double (resp. triple) cD/3 point $P \in C$. If (X, C) satisfies 4.8.3, then (X, C) is an isolated (resp. a divisorial) extremal curve germ.*

Proof. In the hypothesis of 4.1 we additionally assume that $\ell(P) \geq 3$.

4.9. Lemma. *Under notation of 4.8 X has no type (III) points.*

Proof. Assume that X has a type (III) point R . We derive a contradiction. By Lemma 4.7 $\ell(P) = 3$ or 4.

4.9.1. Case $\ell(P) = 3$. We claim that $H^1(\text{gr}_C^2 \omega) \neq 0$. By [Mor88, 2.16] $i_P(1) = 2$ and (in some coordinate system) α satisfies $\alpha \equiv y_1^3 y_4 \pmod{(y_2, y_3, y_4)^2}$ (and $C^\sharp$ is the y_1 -axis). If α contains the term $y_1^k y_2 y_3$, then $k \geq 3$ and this term can be removed by the coordinate change $y_4 \mapsto y_4 - y_1^{k-3} y_2 y_3$. Hence, we may assume that

$$\alpha \equiv y_1^3 y_4 + \lambda y_1 y_2^2 + \mu y_1^2 y_3^2 \pmod{(y_2, y_3)^3 + y_4(y_2, y_3, y_4)} \quad (\subset I_C^{(3)\sharp})$$

for some $\lambda, \mu \in \mathcal{O}_X \pmod{I_C}$. The functions y_2, y_3 form an ℓ -basis of $\text{gr}_C^1 \mathcal{O}$ at P . Since

$$\deg \text{gr}_C^1 \mathcal{O} = 1 - i_P(1) - i_R(1) = -2$$

and $H^1(\text{gr}_C^1 \mathcal{O}) = 0$, we have $\text{gr}_C^1 \mathcal{O} = \mathcal{O}(-1) \oplus \mathcal{O}(-1)$. Furthermore, by [KM92, (2.8)], one has

$$\text{gr}_C^1 \mathcal{O} = (-1 + P^\sharp) \tilde{\oplus} (-1 + 2P^\sharp).$$

where y_3 (resp. y_2) is an ℓ -free ℓ -basis of $(-1 + P^\sharp)$ (resp. of $(-1 + 2P^\sharp)$) at P . Let σ be an ℓ -basis of ω . By the above, one has

$$\omega \tilde{\otimes} \tilde{S}^2 \text{gr}_C^1 \mathcal{O} = (-2 + P^\sharp) \tilde{\oplus} (-2 + 2P^\sharp) \tilde{\oplus} (-1),$$

where $y_3^2 \sigma$ (resp. $y_3 y_2 \sigma, y_2^2 \sigma$) is an ℓ -free ℓ -basis of $(-2 + P^\sharp)$ (resp. $(-2 + 2P^\sharp), (-1)$) at P . There is an injection of coherent sheaves

$$\iota : \omega \tilde{\otimes} \tilde{S}^2 \text{gr}_C^1 \mathcal{O} \longrightarrow \text{gr}_C^2 \omega.$$

As an abstract sheaf, $\omega \tilde{\otimes} \tilde{S}^2 \text{gr}_C^1 \mathcal{O}$ at P is generated by sections $y_3^2 y_1 \sigma, y_3 y_2 y_1^2 \sigma, y_2^2 \sigma$. Further, it is easy to see that $I_C^{(2)\sharp} / I_C^{(3)\sharp}$ at P is generated by elements y_4, y_3^2 ,

y_2y_3, y_2^2 . Hence, $\text{gr}_C^2 \omega$ at P is generated by $y_4y_1^2\sigma, y_3^2y_1\sigma, y_2y_3y_1^2\sigma, y_2^2\sigma$. On the other hand, $y_4 \in I_C^{(2)}$ and $y_1^2y_2y_4, y_1y_3y_4 \in I_C^{(3)}$. By our expression of α

$$(y_1^2y_4 + \lambda y_2^2 + \mu y_1y_3^2)\sigma = 0 \quad \text{in } \text{gr}_C^2 \omega \text{ at } P.$$

Hence, $\text{gr}_C^2 \omega$ at P is generated by the elements $y_3^2y_1\sigma, y_2y_3y_1^2\sigma, y_2^2\sigma$. This means that ι is an isomorphism at P .

Since $i_R(1) = 1$, by [Mor88, 2.16] $\ell(R) = 1$ and in some coordinate system the local equation $\beta(z_1, \dots, z_4) = 0$ of (X, R) satisfies $\beta \equiv z_1z_2 \pmod{(z_2, z_3, z_4)^2}$, where C is the z_1 -axis. Then locally near R we have $I_C^{(2)} = (z_3^2, z_3z_4, z_4^2, z_2)$, so

$$\text{gr}_C^1 \mathcal{O} = \mathcal{O}z_3 \oplus \mathcal{O}z_4 \quad \text{and} \quad S^2 \text{gr}_C^1 \mathcal{O} = \mathcal{O}z_3^2 \oplus \mathcal{O}z_4^2 \oplus \mathcal{O}z_3z_4.$$

Furthermore, $\text{gr}_C^2 \mathcal{O}$ is generated by $z_2, z_3^2, z_4^2, z_3z_4$. Hence, z_2 generates $\text{Coker } \iota$ and so $\text{len}_R \text{Coker } \iota \leq 1$. In this case, $\dim H^0(\text{Coker } \iota) \leq 1$ and $\dim H^1(\text{Im } \iota) = 2$. Therefore, $H^1(\text{gr}_C^2 \omega) \neq 0$ as claimed.

Now from $H^0(\text{gr}_C^j \omega) = 0$, where $j = 0, 1$ and the exact sequences

$$0 \longrightarrow \text{gr}_C^n \omega \longrightarrow \omega_X / F^{n+1} \omega_X \longrightarrow \omega_X / F^n \omega_X \longrightarrow 0, \quad n = 1, 2$$

we have $H^1(\omega_X / F^3 \omega_X) \neq 0$. If f is birational, then by [Mor88, 1.2-1.2.1] we get a contradiction. Assume that f is a $\mathbb{Q}$ -conic bundle. Put $V := \text{Spec}_X \mathcal{O}_X / I_C^{(3)}$. By [MP08, Th. 4.4] $V \supset f^{-1}(o)$. Since

$$-K_X \cdot V = -6K_X \cdot C = 2 = -K_X \cdot f^{-1}(o),$$

we have $V = f^{-1}(o)$. Let $P \in C$ be a general point. Then in a suitable coordinate system (x, y, z) near P we may assume that C is the z -axis. So, $I_C = (x, y)$ and $I_C^{(3)} = (x^3, x^2y, xy^2, y^3)$. But then $V = f^{-1}(o)$ is not a local complete intersection near P , a contradiction. This disproves the case 4.9.1.

4.9.2. Case $\ell(P) = 4$. By deformation $\alpha_\lambda = \alpha + \lambda y_1^3 y_4$ at (X, P) we get a germ (X_λ, C_λ) with a point P_λ of type cD/3 with $\ell(P_\lambda) = 3$ (see Theorem 3.2). Moreover, X_λ has a point R_λ of type (III). This is impossible by 4.9.1.

This proves 4.9. □

From now on we treat the case where P is the only singular point of X and $\ell(P) \geq 3$.

4.10. Lemma (cf. [KM92, Lemma 6.12]). *In the notation of 4.1 we have $\ell(P) \leq 4$ and $i_P(1) \leq 2$.*

Proof. Assume that $\ell(P) \geq 5$. Similarly to [KM92, Lemma 6.12] and Lemma 4.7 we write $\alpha \equiv y_1^{\ell(P)} y_j \pmod{(y_2, y_3, y_4)^2}$, where $j \in \{2, 3, 4\}$ and $\ell(P) + \text{wt } y_j \equiv 0 \pmod{3}$, and use deformation $\alpha_\lambda = \alpha + \lambda y_1^{\ell(P)-3} y_j$ (see Theorem 3.2). We get a germ (X_λ, C_λ) with a type (III) point R_λ and a point P_λ of type cD/3 with $\ell(P_\lambda) = \ell(P) - 3$. If $\ell(P) \geq 6$, we get a contradiction by the case 4.9 considered above.

Hence $\ell(P) = 5$, and $X \setminus \{P\}$ is smooth by Lemma 4.7. Then $\alpha \equiv y_1^5 y_2 \pmod{(y_2, y_3, y_4)^2}$, $\deg \text{gr}_C^1 \mathcal{O}_X = -1$, and y_4, y_3 form an ℓ -basis for $\text{gr}_C^1 \mathcal{O}_X$. Thus H is normal at P by Corollary 4.3, and we see that $\text{gr}_C^1 \mathcal{O}_X = (0) \hat{\oplus} (-1 + P^\sharp)$, H is smooth outside P , y_3 is an ℓ -basis of $\text{gr}_C^1 \mathcal{O}_H$, and $\text{gr}_C^1 \mathcal{O}_H = (-1 + P^\sharp)$. We also see

$$\text{gr}_C^0 \omega_H = \text{gr}_C^0 \omega_X = (-1 + 2P^\sharp) \quad \text{and} \quad \text{gr}_C^1 \omega_H = (-1).$$

We note that $C^\sharp = y_1\text{-axis} \subset H^\sharp \subset \mathbb{C}_{y_1, y_2, y_3}^3$, and $H^\sharp = \{\beta = 0\}$, where

$$\beta \equiv y_1^5 y_2 + c y_1^2 y_3^2 \pmod{(y_2^2, y_2 y_3, y_3^3)},$$

and $c \in \mathbb{C}$. We claim $c \neq 0$. Indeed otherwise we have $y_2 \in \mathcal{O}_H(-3C)^\sharp$, whence $\text{gr}_C^2 \mathcal{O}_H^\sharp = \mathcal{O}_{C^\sharp} y_3^2$ and $\text{gr}_C^2 \mathcal{O}_H = (\text{gr}_C^1 \mathcal{O}_H)^{\tilde{\otimes} 2} = (-2 + 2P^\sharp)$. Thus $H^1(H, \mathcal{O}_H) \neq 0$, a contradiction. Hence $c \neq 0$.

Since P is a cD/3 point, we have $y_2 y_3 \notin \alpha$ and $y_3^3 \in \alpha$, and hence $y_2 y_3 \notin \beta$ and $y_3^3 \in \beta$. Since $c \neq 0$, the terms $\gamma(y_1) y_1^3 y_2 y_3$ can be killed by a μ_3 -coordinate change $y_3 \mapsto y_3 - \gamma(y_1) y_1 y_2 / (2c)$ and we may further assume

$$(4.10.1) \quad \beta \equiv y_1^5 y_2 + c y_1^2 y_3^2 + y_3^3 \pmod{(y_2^2, y_2 y_3, y_3^4)}.$$

We claim that $\text{gr}_C^2 \mathcal{O}_H = (-1 + 2P^\sharp)$ and $\text{gr}_C^3 \mathcal{O}_H = (-1)$. First by $y_2 \in \mathcal{O}_H(-2C)^\sharp$, one has $y_1^2(y_1^3 y_2 + c y_3^2) \in \mathcal{O}_H(-3C)^\sharp$. Hence if we set $z := y_1^3 y_2 + c y_3^2$, then $z \in \mathcal{O}_H(-3C)^\sharp$ and $y_3^2 \equiv -y_1^3 y_2 / c \pmod{(z)}$. Thus by $\mathcal{O}_H(-2C)^\sharp = (y_2, y_3^2)$, we see

$$\mathcal{O}_H(-2C)^\sharp / (y_2^2, y_2 y_3, z) = \mathcal{O}_{C^\sharp} y_2 \simeq \mathcal{O}_{C^\sharp} \quad \text{and} \quad \mathcal{O}_H(-3C)^\sharp = (y_2^2, y_2 y_3, z).$$

We also have $y_1^2 z + y_3^3 \in (y_2^2, y_2 y_3^2, y_3^4)$ by (4.10.1), whence $z \equiv y_1 y_2 y_3 / c \pmod{(y_2^2, y_2 y_3^2, z y_3)}$. Thus

$$\mathcal{O}_H(-3C)^\sharp / (y_2^2, y_2 y_3^2, z y_3) = \mathcal{O}_{C^\sharp} y_2 y_3 \simeq \mathcal{O}_{C^\sharp} \quad \text{and} \quad \mathcal{O}_H(-4C)^\sharp = (y_2^2, y_2 y_3^2, z y_3).$$

From these follows the claim:

$$\text{gr}_C^2 \mathcal{O}_H = (\text{gr}_C^1 \mathcal{O}_H)^{\tilde{\otimes} 2} (3P^\sharp) = (-1 + 2P^\sharp)$$

and

$$\text{gr}_C^3 \mathcal{O}_H = \text{gr}_C^1 \mathcal{O}_H \tilde{\otimes} \text{gr}_C^2 \mathcal{O}_H = (-1).$$

We then claim $H^1(\omega_H / \omega_H(-4C)) \neq 0$. Indeed this follows from

$$\text{gr}_C^2 \omega_H = \text{gr}_C^0 \omega_H \tilde{\otimes} \text{gr}_C^2 \mathcal{O}_H = (-1 + P^\sharp)$$

and

$$\text{gr}_C^3 \omega_H = \text{gr}_C^0 \omega_H \tilde{\otimes} \text{gr}_C^3 \mathcal{O}_H = (-2 + 2P^\sharp).$$

Since $\omega_H = \omega_X \tilde{\otimes} \mathcal{O}_H$, the non-vanishing $H^1(\omega_H / \omega_H(-4C)) \neq 0$ means that f is a $\mathbb{Q}$ -conic bundle [Mor88, Remark 1.2.1] and the subscheme $4C$ of H contains the scheme-theoretic fiber $f^{-1}(o)$ [MP08, Th. 4.4]. However

$$-K_X \cdot 4C = 4/3 < 2 = -K_X \cdot f^{-1}(o),$$

a contradiction. The case $\ell(P) = 5$ is thus disproved. $\square$

4.11. Case $\ell(P) = 3$ and no type (III) points. By [Mor88, 2.16] $i_P(1) = 2$ and (in some coordinate system) α satisfies $\alpha \equiv y_1^3 y_4 \pmod{(y_2, y_3, y_4)^2}$ (and $C^\sharp$ is the y_1 -axis). Hence, y_2, y_3 form an ℓ -basis of $\text{gr}_C^1 \mathcal{O}$. Since $\deg \text{gr}_C^1 \mathcal{O} = 1 - i_P(1) = -1$ and $H^1(\text{gr}_C^1 \mathcal{O}) = 0$, $\text{gr}_C^1 \mathcal{O} = \mathcal{O} \oplus \mathcal{O}(-1)$. Further, by [KM92, (2.8)] there are only two possibilities:

$$\text{gr}_C^1 \mathcal{O} = \begin{cases} (2P^\sharp) & \tilde{\oplus} (-1 + P^\sharp), \\ (P^\sharp) & \tilde{\oplus} (-1 + 2P^\sharp). \end{cases}$$

Consider the first case, i.e., $\text{gr}_C^1 \mathcal{O} = (2P^\sharp) \tilde{\oplus} (-1 + P^\sharp)$. Then the arguments in the first part of proof of [KM92, (6.13)] can be applied. Let J be the C -laminal ideal of width 2 such that $J / F_C^2 \mathcal{O} = (2P^\sharp)$. Then we conclude that $H^1(\omega / F^4(\omega, J)) \neq 0$ [KM92, p. 599-600]. If the contraction f is birational, we get a contradiction by

[Mor88, 1.2-1.2.1]. Let f be a $\mathbb{Q}$ -conic bundle. Put $V := \operatorname{Spec}_X \mathcal{O}_X / F^4(\mathcal{O}, J)$. Then $V \equiv mC$ for some m . By [MP08, Th. 4.4] $V \supset f^{-1}(o)$. Hence, $m/3 = -K_X \cdot V < 2 = -K_X \cdot f^{-1}(o)$. On the other hand, near a general point $S \in C$, J is generated by (z_2, z_3^2) , where (z_1, z_2, z_3) are some local coordinates such that C is the z_1 -axis. Hence, $F^4(\omega, J) = J^2 = (z_2, z_3^2)^2$ near S . So, $m = \operatorname{len}(\mathbb{C}[z_2, z_3]/F^4(\omega, J)) = 6$. Therefore, $f^{-1}(o) = V$ and its ideal sheaf coincides with $F^4(\omega, J)$. However, $F^4(\omega, J)$ is not generated by two elements near S , so $f^{-1}(o)$ is not a locally complete intersection, a contradiction.

Consider the second case, i.e., $\operatorname{gr}_C^1 \mathcal{O} = (P^\#) \tilde{\oplus} (-1 + 2P^\#)$. If (X, P) is a double cD/3 point, then f is a flipping contraction by [KM92, Theorem (6.3)], whence we get the configuration (4.8.5.1). Thus we assume that the term $y_1 y_2^2$ does not appear in α . Further we use arguments of the proof of [KM92, Lemma (6.13), p. 600]. Let J be the C -laminal ideal of width 2 such that $J/F_C^2 \mathcal{O} = (P^\#)$. Modulo a μ_3 -equivariant change of coordinates we may further assume that y_3 (resp. y_2) is an ℓ -free ℓ -basis of $(P^\#)$ (resp. $(-1 + 2P^\#)$) in $\operatorname{gr}_C^1 \mathcal{O}$ and that $\alpha \equiv y_1^3 y_4 \pmod{I^\# J^\#}$. Whence $J^\# = (y_2^2, y_3, y_4)$ at $P^\#$ and $y_4 \in F^3(\mathcal{O}, J)^\#$. Let K be the ideal such that $J \supset K \supset F^3(\mathcal{O}, J)$ and $K/F^3(\mathcal{O}, J) = (P^\#)$ in

$$\operatorname{gr}^2(\mathcal{O}, J) = \operatorname{gr}^{2,0}(\mathcal{O}, J) \tilde{\oplus} \operatorname{gr}^{2,1}(\mathcal{O}, J) = (P^\#) \tilde{\oplus} (-1 + P^\#).$$

Here we may assume that y_3 (resp. y_2^2) is an ℓ -free ℓ -basis of $(P^\#)$ (resp. $(-1 + P^\#)$) in the above ℓ -splitting modulo a coordinate change $y_3 \mapsto y_3 + (\cdots)y_2^2$. We then have $K^\# = (y_2^3, y_3, y_4)$ at $P^\#$ and

$$\operatorname{gr}^1(\mathcal{O}, K) = (-1 + 2P^\#), \quad \operatorname{gr}^2(\mathcal{O}, K) = (-1 + P^\#).$$

We have $\operatorname{gr}^{3,0}(\mathcal{O}, K) \simeq \operatorname{gr}^{2,0}(\mathcal{O}, J) \simeq (P^\#)$ and

$$\alpha \equiv y_1^3 y_4 + c y_2^3 \pmod{I^\# K^\#},$$

for some unit $c \in \mathcal{O}_X^\times$, because $I^\# J^\# = I^\# K^\# + (y_2^3)$ and $y_2^3 \in \alpha$. Whence we have an ℓ -isomorphism

$$\operatorname{gr}^{3,1}(\mathcal{O}, K) \simeq \operatorname{gr}^1(\mathcal{O}, K)^{\tilde{\otimes} 3} (3P^\#) \simeq (0)$$

as in [KM92, p. 600], and an ℓ -splitting

$$\operatorname{gr}^3(\mathcal{O}, K) = \operatorname{gr}^{3,0}(\mathcal{O}, K) \tilde{\oplus} \operatorname{gr}^{3,1}(\mathcal{O}, K),$$

in which y_3 (resp. y_4) is an ℓ -free ℓ -basis of $(P^\#)$ (resp. (0)) modulo a coordinate change $y_3 \mapsto y_3 + (\cdots)y_1^2 y_4$. For any $l > 0$ there is a natural exact sequence

$$(4.11.1) \quad 0 \longrightarrow F^{l+1}(\mathcal{O}, K) \longrightarrow F^l(\mathcal{O}, K) \longrightarrow \operatorname{gr}^l(\mathcal{O}, K) \longrightarrow 0.$$

We claim that the sections $y_1 y_3, y_4 \in \operatorname{gr}^3(\mathcal{O}, K)$ can be extended to sections of $F^3(\mathcal{O}, K) = F^1(K)$. By (4.11.1) it is sufficient to show that $H^1(F^4(\mathcal{O}, K)) = 0$. There are injections of coherent sheaves

$$\begin{aligned} \operatorname{gr}^{3n}(\mathcal{O}, K) &\hookleftarrow \tilde{S}^n \operatorname{gr}^3(K), \\ \operatorname{gr}^{3n+1}(\mathcal{O}, K) &\hookleftarrow \tilde{S}^n \operatorname{gr}^3(K) \tilde{\otimes} \operatorname{gr}^1(\mathcal{O}, K), \\ \operatorname{gr}^{3n+2}(\mathcal{O}, K) &\hookleftarrow \tilde{S}^n \operatorname{gr}^3(K) \tilde{\otimes} \operatorname{gr}^2(\mathcal{O}, K) \end{aligned}$$

with cokernels of finite length. Therefore, for any $l > 0$, the degree of each component in a decomposition of $\operatorname{gr}^l(\mathcal{O}, K)$ in a direct sum is at least -1 . Then $H^1(\operatorname{gr}^l(\mathcal{O}, K)) = 0$ and from (4.11.1) we get surjections

$$H^1(F^{l+n}(\mathcal{O}, K)) \rightarrow H^1(F^l(\mathcal{O}, K)) \quad \text{for } l, n > 0.$$

Hence, $H^1(F^l(\mathcal{O}, K)/F^{l+n}(\mathcal{O}, K)) = 0$. Note that for any $m > 0$ there is $n > 0$ such that $I_C^m F^l(\mathcal{O}, K) \supset F^{l+n}(\mathcal{O}, K)$. By the Formal Function Theorem we have

$$\begin{aligned} H^1(F^l(\mathcal{O}, K))^\wedge &= H^1(F^l(\hat{\mathcal{O}}, K)) = \varprojlim H^1(F^l(\mathcal{O}, K)/I_C^m F^l(\mathcal{O}, K)) = \\ &= \varprojlim H^1(F^l(\mathcal{O}, K)/F^{l+n}(\mathcal{O}, K)) = 0. \end{aligned}$$

Hence $H^1(F^l(\mathcal{O}, K)) = 0$ for $l > 0$ and there are surjections

$$H^0(F^l(\mathcal{O}, K)) \longrightarrow H^0(\mathrm{gr}^l(\mathcal{O}, K)) \longrightarrow 0.$$

This proves our claim. Therefore, near P a general member $H \in |\mathcal{O}_X|_C$ is given by equations $\alpha(y_1, \dots, y_4) = 0$ and $\beta(y_1, \dots, y_4) = 0$, where $\alpha = y_4^2 + y_3^3 + y_2^3 +$ (terms of degree ≥ 4) (recall that $\alpha \not\equiv y_1^2 y_2, y_1 y_2^2$), $\beta \equiv \lambda y_3 y_1 + y_4 \pmod{F^4(\mathcal{O}, K)}$, and $\lambda \in \mathcal{O}_{\mathbb{C}^4}$ such that $\lambda(P) \in \mathbb{C}$ can be chosen arbitrarily. Hence we can eliminate y_4 and get


$$(H, P) = \{\gamma(y_1, y_2, y_3) = 0\}/\mu_3(1, 1, 2) \supset C = y_1\text{-axis}/\mu_3,$$

where γ is a μ_3 -invariant convergent power series such that, for $\sigma = (1, 1, 2)$, $\gamma_{\sigma=3} = y_2^3$ and the term $\gamma_{\sigma=6}(y_1, 0, y_3)$ is squarefree. Hence we are done by Computation 4.12 below.

4.12. Computation. Let (D, P) be a normal surface singularity

$$(D, P) = \{\gamma = 0\}/\mu_3 \subset \mathbb{C}^3/\mu_3(1, 1, 2),$$

where $\gamma = \gamma(y_1, y_2, y_3)$ is μ_3 -invariant and let $C := (y_1\text{-axis})/\mu_3$. Let σ be the weight $(1, 1, 2)$. Assume that $\gamma_{\sigma=3} = y_2^3$ and $\gamma_{\sigma=6}(y_1, 0, y_3)$ is squarefree. Then D has only rational singularities and $\Delta(D, C)$ is as follows.



Sketch of the proof. We note that $\gamma_{\sigma=6}(y_1, 0, y_3)$ contains y_3^3 since it is squarefree. Consider the weighted blowup $\hat{H} \rightarrow H$ with weights $\frac{1}{3}(1, 1, 2)$. The exceptional divisor Λ is given by $\gamma_{\sigma=3} = y_2^3 = 0$ in the weighted projective plane $\mathbb{P}(1, 1, 2)$. Hence Λ is a smooth rational curve. Clearly, $\mathrm{Sing}(\hat{H})$ is contained in Λ . In the chart $U_1 := \{y_1 \neq 0\}$ the surface $\hat{H}$ is given by

$$y_2^3 + y_1 \gamma_{\sigma=6}(1, y_2, y_3) + y_1^2 \gamma_{\sigma=9}(1, y_2, y_3) + \dots = 0.$$

Hence, $\mathrm{Sing}(\hat{H}) \cap U_1$ is given by $y_1 = y_2 = \gamma_{\sigma=6}(1, 0, y_3) = 0$. Since $\gamma_{\sigma=6}(1, 0, y_3)$ is a cubic polynomial without multiple factors, $\mathrm{Sing}(\hat{H}) \cap U_1$ consists of three points: $P_0 := (0, 0, 0)$, P_1 , P_2 . In particular, this shows that $\hat{H}$ is normal. Further, $\gamma_{\sigma=6}(1, y_2, y_3)$ contains the term y_3 . Hence at the origin $\hat{H}$ has a Du Val singularity of type A_2 and the pair

$$(\hat{H}, \Lambda + \hat{C}) \simeq (\{y_2^3 + y_1 y_3 = 0\}, \{y_2 = 0\})$$

is LC, where $\hat{C}$ is the proper transform of C . This gives us the left-hand side of the graph. Similarly, from P_1 and P_2 we get the upper and the right-hand side of the graph. The vertex $\circ$ in the bottom comes from the chart $y_3 \neq 0$. The computation of the self-intersection number of the central vertex is an easy exercise. $\square$

4.13. Case $\ell(P) = 4$ and no type (III) points. By [Mor88, 2.16] $i_P(1) = 2$ and (in some coordinate system) α satisfies $\alpha \equiv y_1^4 y_3 \pmod{(y_2, y_3, y_4)^2}$ (and $C^\#$ is the y_1 -axis). Hence, y_2, y_4 form an ℓ -basis of $\text{gr}_C^1 \mathcal{O}$.

We prove Claim 4.8.3. Since it has been proved that a type (III) point does not occur, it remains to settle the ℓ -isomorphism (4.8.3.1). If it does not hold, then we have $\text{gr}_C^1 \mathcal{O} = (2P^\#) \tilde{\oplus} (-1)$ and $\text{gr}_C^1 \omega = (P^\#) \tilde{\oplus} (-2 + 2P^\#)$, whence $H^1(\text{gr}_C^1 \omega) \neq 0$. Thus we get a contradiction as in 4.11 and Claim 4.8.3 is proved.

If (X, C) is flipping, then Claims 4.8.2, 4.8.4, and 4.8.5 are already proved in [KM92, (6.3)]. Since $\ell(P) > 2$, P is a double or triple cD/3 point, and Claim 4.8.6 is proved in [KM92, (6.3.4)] if P is a double cD/3 point.

Assume that (X, C) is not isolated. Then P , as a cD/3 point, is triple by $\ell(P) > 2$ and [KM92, (6.3.4)]. This proves 4.8.2.

Let J be the C -laminal ideal of width 2 such that $J/F_C^2 \mathcal{O} = (0)$ in the ℓ -splitting (4.8.3.1). Up to coordinate change we may assume that y_4 (resp. y_2) is an ℓ -free ℓ -basis of (0) (resp. $(-1 + 2P^\#)$) in $\text{gr}_C^1 \mathcal{O}$ and that $\alpha \equiv y_1^4 y_3 \pmod{I_C^\# J^\#}$. Whence $y_3 \in F^3(\mathcal{O}, J)^\#$. We note that $y_1 y_2^2 \notin \alpha$ in the new coordinates since P is a triple cD/3 point.

Since we have ℓ -isomorphisms

$$\begin{aligned} \text{gr}^{2,0}(\mathcal{O}, J) &\simeq \text{gr}^0(\mathcal{O}, J) \simeq (0) \\ \text{gr}^{2,1}(\mathcal{O}, J) &\simeq \text{gr}^1(\mathcal{O}, J)^{\otimes 2} \simeq (-1 + P^\#), \end{aligned}$$

the ℓ -exact sequence

$$0 \rightarrow \text{gr}^{2,1}(\mathcal{O}, J) \rightarrow \text{gr}^2(\mathcal{O}, J) \rightarrow \text{gr}^{2,0}(\mathcal{O}, J) \rightarrow 0,$$

is ℓ -split. Let K be the ideal such that $J \supset K \supset F^3(\mathcal{O}, J)$ and $K/F^3(\mathcal{O}, J) = (0)$ in

$$\text{gr}^2(\mathcal{O}, J) \simeq (0) \tilde{\oplus} (-1 + P^\#).$$

Here we may assume that y_4 (resp. y_2^2) is an ℓ -free ℓ -basis of (0) (resp. $(-1 + P^\#)$) modulo a coordinate change $y_4 \mapsto y_4 + (\cdots) y_1 y_2^2$.

We have thus $K^\# = (y_2^3, y_3, y_4)$ and

$$\text{gr}^1(\mathcal{O}, K) = (-1 + 2P^\#), \quad \text{gr}^2(\mathcal{O}, K) = (-1 + P^\#).$$

We have $\text{gr}^{3,0}(\mathcal{O}, K) \simeq \text{gr}^{2,0}(\mathcal{O}, J) \simeq (0)$ and

$$\alpha \equiv y_1^4 y_3 + c y_2^3 \pmod{I^\# K^\#},$$

for some unit $c \in \mathcal{O}_X^\times$, because $I^\# J^\# = I^\# K^\# + (y_2^3)$ and $y_2^3 \in \alpha$. Whence we have an ℓ -isomorphism

$$\text{gr}^{3,1}(\mathcal{O}, K) \simeq \text{gr}^1(\mathcal{O}, K)^{\otimes 3} (4P^\#) \simeq (P^\#).$$

Thus we have an ℓ -splitting

$$\text{gr}^3(\mathcal{O}, K) \simeq \text{gr}^{3,0}(\mathcal{O}, K) \tilde{\oplus} \text{gr}^{3,1}(\mathcal{O}, K) \simeq (0) \tilde{\oplus} (P^\#).$$

By a change of coordinate $y_4 \mapsto y_4 + (\cdots) y_1 y_3$, we may further assume that y_4 (resp. y_3) is an ℓ -free ℓ -basis of (0) (resp. $(P^\#)$). By the same computation as in 4.11, we get the configuration (4.8.5.2). This contracts to a Du Val point of type E_6 , and hence f is a divisorial contraction, which proves 4.8.1.

Finally we note that [KM92, (6.15) and (6.20)] settled 4.8.4 for isolated (X, C) , and 4.8.6 for a double $cD/3$ point. We omit the proofs of 4.8.4 and 4.8.6 in other cases since the arguments are similar. This completes our treatment of the case $\ell(P) > 2$. $\square$

4.14. Examples. To show that all the possibilities 1.4.1, 1.4.2, and 1.4.3 occur we use deformation arguments. Consider the surface contraction $f_H : H \rightarrow T$ with dual graph of the form 1.4.1 or 1.4.2. By [KM92, Proposition (11.4)] the natural map from the deformation space of H to the product of deformation spaces of singularities $P, R \in H$ is smooth, in particular, surjective. Moreover, the total deformation space $\mathfrak{X}$ of H has a morphism $\mathfrak{f}$ to the total deformation space $\mathfrak{X}_Z$ of T so that $\mathfrak{f}|_H = f_H$. This means in particular that any $\mathbb{Q}$ -Gorenstein deformation of singularities of H can be globalized. Now assume that (H, P) and (H, R) can be obtained as hyperplane sections of some terminal singularities (X, P) and (X, R) respectively. Regard (X, P) and (X, R) as deformation spaces of (H, P) and (H, R) respectively. By the above there is a globalization $f : X \supset H \rightarrow Z \supset T$.

4.14.1. Example. Consider the surface contraction $f_H : H \rightarrow T$ with dual graph 1.4.1 and consider the following terminal singularities:

$$\begin{aligned} (X, P) &= \{y_4^2 + y_3^3 + y_1 y_2 (y_1 + y_2) = 0\} / \mu_3(1, 1, 2, 0), \\ (X, R) &= \{z_1 z_2 + z_3^2 + z_4^m = 0\}, \quad m \geq 1. \end{aligned}$$

Let $H \subset (X, P)$ is given by $y_4 = 0$ and $H \subset (X, R)$ is given by $z_4 = 0$. By [KM92, (6.7.1)] the dual graph of the minimal resolution of (H, P) is the same as that in 1.4.1. By 4.14 one obtains the corresponding birational contraction $f : X \supset H \rightarrow Z \supset T$. Here (X, P) is a simple $cD/3$ -singularity (see [Rei87]). Therefore, this f is a divisorial contraction of type 1.4.1. The point $R \in X$ is smooth if $m = 1$ and is a cA_1 -singularity if $m > 1$.

4.14.2. Example. Similarly to Example 4.14.1, take

$$(X, P) = \{y_4^2 + y_1^2 y_2 + y_2^6 + y_3^3 = 0\} / \mu_3(1, 1, 2, 0).$$

By [KM92, (6.7.2)] we get an example of a divisorial contraction as in 1.4.2.

4.14.3. Example. As above, take

$$(X, P) = \{y_2^3 + y_3^3 + y_3 y_1^4 + y_4^2\} / \mu_3(1, 1, 2, 0),$$

where H is cut out by $y_4 = y_1 y_3$. We get an example of a divisorial contraction as in 1.4.3.

5. CASE: P IS OF TYPE cA/m AND H IS NORMAL

In this section we prove Theorems 1.5 and 1.8 in the case where a general $H \in |\mathcal{O}_X|_C$ is normal. Thus throughout this section we assume that (X, C) is an extremal curve germ of type (IA) or $(IA)^\vee$ such that the only non-Gorenstein point $P \in X$ is of type cA/m (see 1.7). Let $F \in |-K_X|$ be a general member. Take $H \in |\mathcal{O}_X|_C$ so that the pair $(X, F + H)$ is LC (see Proposition 2.6). Assume that H is normal. Let $f : (X, C) \rightarrow (Z, o)$ be the corresponding contraction.

5.1. Proposition. *In the above notation, H has only log terminal singularities of type T. Furthermore, the pair (H, C) is PLT outside of P and $H \setminus \{P\}$ has at most one singular point, which if exists is Du Val of type A_n . If moreover f is birational,*


then $\Delta(H, C)$ is as in (1.8.1.1). If moreover f is a $\mathbb{Q}$ -conic bundle, then $\Delta(H, C)$ is of the form

$$(5.1.1) \quad \circ - \circ - \circ - \bullet - \overset{4}{\circ}$$

In particular, $m = 2$ and (X, P) is either a cyclic quotient singularity $\frac{1}{2}(1, 1, 1)$ or a singularity of the form $\{xy + z^2 + t^k = 0\}/\mu_2(1, 1, 1, 0)$.

Proof. First we claim that H has only log terminal singularities. Write $K_H + F|_H = (K_X + H + F)|_H \sim 0$. Recall that $F \cap C = \{P\}$. So $(H, F|_H)$ is not klt at P and klt at a general point of C . We see that $(H, F|_H)$ is klt outside of P by the Connectedness Lemma [Sho93, 5.7], [Kol92, 17.4] (if f is birational) and by Lemma 2.3.2 (if f is a $\mathbb{Q}$ -conic bundle). On the other hand, by our assumptions and the adjunction formula the pair $(H, F|_H)$ is LC near $F \cap H$, so the surface H has at worst log terminal singularities. Further, since H is a Cartier divisor in X , the singularities of H are of type T (see 2.2.1).

Now we claim that the pair (H, C) is PLT outside of P . Assume that $K_H + C$ is not PLT at some point $Q \neq P$. Take c so that $(H, F|_H + cC)$ is maximally LC. By the Connectedness Lemma and Lemma 2.3.2 we have $c = 1$, so $(H, F|_H + C)$ is LC. Therefore, H has a log terminal singularity at Q and the point (H, Q) is Du Val. From the classification of log canonical pairs (see, e.g., [Kol92, Ch. 3]) we obtain that the part of the dual graph $\Delta(H, C)$ which represents H near the singularity Q is of the form



But then the corresponding matrix of this subgraph is not negative definite, a contradiction. Thus (H, C) is PLT outside of P . Since any point $Q \in H \setminus \{P\}$ is Gorenstein, it is Du Val of type A_n or smooth. Near each such a point the dual graph $\Delta(H, C)$ is of the form



If (H, C) contains two such points, we get a contradiction with negative definiteness of the corresponding matrix. Thus we obtain (1.8.1.1).

Now consider the case where f is a $\mathbb{Q}$ -conic bundle. If (H, C) is PLT also at P , then H has two singularities of types $\frac{1}{n}(1, q)$ and $\frac{1}{n}(1, n - q)$ (see Lemma 2.3.1). Since they are of type T, we see the following by Proposition 2.2.2:

$$(q + 1)^2 \equiv 0 \pmod{n}, \quad (n - q + 1)^2 \equiv 0 \pmod{n}.$$

This gives us $4 \equiv 0 \pmod{n}$. Since X is not Gorenstein, the singularities of H are worse than Du Val. Hence, $n = 4$. We get the graph (5.1.1).

Finally assume that (H, C) is not PLT at P . Then $\Delta(H, C)$ is of the form (1.8.1.1) with $r \neq 1$, $r \neq n$, and $c_1 c_n \geq 6$ by Proposition 2.2.3. Contracting black vertices successively, on some step we get a subgraph

$$(5.1.2) \quad \overset{c_1}{\circ} - \dots - \overset{c_{r-1}}{\circ} - \bullet - \overset{c_{r+1}}{\circ} - \dots - \overset{c_n}{\circ}$$

Hence strings $[c_{r-1}, \dots, c_1]$ and $[c_{r+1}, \dots, c_n]$ are conjugate. This contradicts the following claim because $c_1 c_n \geq 6$. $\square$

5.1.3. Claim. Let $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate strings. If, for some $c \geq 2$, the string of the form

$$(5.1.3.1) \quad [a_r, \dots, a_1, c, b_1, \dots, b_s]$$


is of type T, then it is Du Val.

Proof. Assume that the string ((5.1.3.1)) is not Du Val. Take it so that $r + s$ is minimal. Since $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate, either $a_r = 2$ or $b_s = 2$. Assume that $a_r = 2$. If $r = 1$, then $s = 1$ and $b_1 = 2$, which is a contradiction by Proposition 2.2.3, (iii). Hence, $r > 1$, $b_s > 2$, and $[a_{r-1}, \dots, a_1, c, b_1, \dots, b_{s-1}, b_s - 1]$ is again a non-Du Val T-string (see Proposition 2.2.2) and the strings $[a_1, \dots, a_{r-1}]$ and $[b_1, \dots, b_{s-1}, b_s - 1]$ are conjugate. This contradicts our minimality assumption. $\square$

Thus (i) of Theorem 1.6 exhausts all $\mathbb{Q}$ -conic bundles with normal H . Explicit examples will be given in Sect. 7.

5.2. In the birational case, similarly to 4.14 any $\mathbb{Q}$ -Gorenstein deformations of singular points of H can be globalized by [KM92, Proposition (11.4)].

5.2.1. Example. Let $[b_1, \dots, b_r]$ be any T-string and let $b_l > 2$. Then the configuration



where $k \leq b_l - 3$, determines a surface germ (H, C) which is contracted to (T, o) with the dual graph

$$b_1 \text{ --- } \dots \text{ --- } b_l - k - 1 \text{ --- } \dots \text{ --- } b_r$$


For example, for $[b_1, \dots, b_r] = [4]$ and $k = 0$, this gives Francia's flip (see [KM92, Theorem 4.7]). For $[b_1, \dots, b_r] = [3, 2, \dots, 2, 3]$, $l = r$, and $k = 1$ this gives examples of divisorial extremal neighborhood of index two [KM92, 4.7.3.1.1].

6. CASE: P IS OF TYPE cA/m AND H IS NOT NORMAL

6.1. In this section we prove Theorems 1.5 and 1.8 in the case where a general $H \in |\mathcal{O}_X|_C$ is not normal. Thus throughout this section we assume that (X, C) is an extremal curve germ of type (IA) or $(IA^\vee)$, the only non-Gorenstein point $P \in X$ is of type cA/m . Let $F \in |-K_X|$ be a general member. Let $H \in |\mathcal{O}_X|_C$ be a non-normal member such that the pair $(X, H + F)$ is LC (see Proposition 2.6). Let $f : (X, C) \rightarrow (Z, o)$ be the corresponding contraction.

6.2. Setup. Let $\nu : H' \rightarrow H$ be the normalization and let $\mu : \tilde{H} \rightarrow H'$ be the minimal resolution. Let $C' = \nu^{-1}(C)$ (with reduced structure) and let $\tilde{C} \subset \tilde{H}$ be the proper transform of C' . If C' is reducible, components of C' (resp. $\tilde{C}$) are denoted by C'_i (resp. $\tilde{C}_i$). Let $\bar{H}$ be a minimal model over T (so that $\bar{H}$ is smooth

and has no (-1) -curves on fibers over T). Thus we have the following diagram:



Let $\Upsilon := \nu^{-1}(F \cap H)$. By 6.1 and Corollary 2.6.1 we have

6.2.1. Corollary. *The pair $(H', C' + \Upsilon)$ is LC and the restriction map $\nu|_{C'} : C' \rightarrow C$ is of degree 2.*

6.2.2. Corollary. *Pull-back $C^\sharp$ of C to the index-one cover $(X^\sharp, P^\sharp) \rightarrow (X, P)$ is smooth. In particular, (X, C) is of type (IA).*

Note that $\Delta(H', C')$ is the dual graph of the 1-cycle $v^{-1}(o) \subset \tilde{H}$. Hence $\Delta(H', C')$ is negative semi-definite and its *fundamental cycle* is defined as usual.

6.3. Proposition. *Under the assumptions of 6.1 the following are equivalent:*

- (i) *every member of $|\mathcal{O}_X|_C$ is non-normal,*
- (ii) *each component of $\tilde{C}$ appears with coefficient > 1 in the fundamental cycle G of $\Delta(H', C')$.*

In particular, if every member of $|\mathcal{O}_X|_C$ is non-normal, then all the components of $\tilde{C}$ are contracted by $\tilde{v} : \tilde{H} \rightarrow \bar{H}$.

Proof. Assume that (ii) does not hold, that is, a component $\tilde{C}_1 \subset \tilde{C}$ appears with coefficient 1 in G . Then there is a function $\psi \in \mathfrak{m}_{o,T}$ such that $v^*\psi$ has a simple zero along $\tilde{C}_1$. Note that the map $H^0(Z, \mathcal{O}_Z) \rightarrow H^0(T, \mathcal{O}_T)$ is surjective. Hence $\psi = \phi|_T$ for some $\phi \in \mathcal{O}_Z$. Pick a general point $S \in C$. If $f^*\phi = 0$ is singular along C , then $f^*\phi \in I_C^2$ at S . By commutativity of the above diagram, we have $v^*\psi = \mu^*\nu^*(f^*\phi)|_H \in I_{\tilde{C}_1}^2$ at a point above S . This contradicts the construction of ψ . So $f^*\phi = 0$ is smooth along C and a general member of $|\mathcal{O}_X|_C$ is normal, so (i) does not hold.

Conversely, assume that (i) does not hold. Then there is a normal member $L \in |\mathcal{O}_X|_C$. Regard X as analytic neighborhood of a general point $Q \in C$. Then $H = H_1 + H_2$, where H_1, H_2 are smooth surfaces intersecting transversely along C . Hence L intersects transversely at least one of H_1, H_2 along C . This means that $\nu^*L|_H$ is reduced along at least one component of C' . Thus (ii) does not hold.


As for the last statement, we note that (T, o) is either a cyclic quotient singularity (see Proposition 2.6) or a smooth curve. In both cases $\tilde{v}(G)$ is reduced. $\square$

6.4. Proposition. *Under the assumptions of 6.1, there are only two possibilities for the dual graph $\Delta(H', C' + \Upsilon)$:*

6.4.1. C' has two irreducible components: $C' = C'_1 + C'_2$.

$$\square - \underbrace{\overset{a_r}{\circ} \cdots \overset{a_1}{\circ}}_{\Delta_1} - \diamond - \underbrace{\overset{c_1}{\circ} \cdots \overset{c_l}{\circ}}_{\Delta_3} - \diamond - \underbrace{\overset{b_1}{\circ} \cdots \overset{b_s}{\circ}}_{\Delta_2} - \square$$

6.4.2. C' is irreducible.




Here $\square$ corresponds to an irreducible component of Υ , $\diamond$ corresponds to an irreducible component of C' , the chain Δ_1 (resp., Δ_2) corresponds to the singularity of type $\frac{1}{m}(1, a)$ (resp., $\frac{1}{m}(1, -a)$), and in case 6.4.1 the chain Δ_3 corresponds to the point (H', Q') , where $Q' = C'_1 \cap C'_2$. The strings $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate. If f is birational, then at least one of the vertices $\diamond$ corresponds to a (-1) -curve under the extra assumption that every member of $|\mathcal{O}_X|_C$ is non-normal. If f is a $\mathbb{Q}$ -conic bundle, then all the vertices $\diamond$ correspond to (-1) -curves.

Proof. Note that C' is a fiber of a contraction $H' \rightarrow T \ni o$, where (T, o) is either a cyclic quotient singularity (see Lemma 2.6) or a curve germ. Hence $p_a(C') = 0$ and all components of C' are smooth rational curves. By Corollary 6.2.1 C' has at most two components. So either $C' \simeq \mathbb{P}^1$ or C' is a union of two $\mathbb{P}^1$'s meeting each other at one point, say Q' .

By the classification of log canonical pairs (see, e.g., [Kol92, Ch. 3]) Υ is smooth at any point $\Upsilon \cap C'$. On the other hand, $\Upsilon = \nu^{-1}(F \cap H)$, where H is Cartier and the pair $(F, H \cap F)$ is LC. Hence Υ has exactly two components Υ_1, Υ_2 and these components are smooth.

Further, since $(H', \Upsilon + C')$ is LC, through any point of H' pass at most two components of $\Upsilon + C'$. Thus for the configuration of $\Upsilon + C'$ on H' we have only the following two possibilities:



Since the pair $(H', \Upsilon + C')$ is LC, from the classification of log canonical pairs (see, e.g., [Kol92, Ch. 3]) we get the desired graphs 6.4.1 and 6.4.2.

It remains to prove the last statements about (-1) -curves. If f is birational, then by Proposition 6.3 at least one of the components of C' is a (-1) -curve. Assume that f is a $\mathbb{Q}$ -conic bundle. Clearly, the fiber $\nu^{-1}(o)$ of a rational curve fibration ν contains a (-1) -curve and this curve must coincide with a component of C' . So we are done if C' is irreducible. Consider the case 6.4.1. By the above one of the $\diamond$ -vertices corresponds to a (-1) -curve. Hence the chain $\Delta_1 - \bullet - \Delta_3 - \diamond - \Delta_2$ forms a fiber of a rational curve fibration and we may assume that $\bullet$ is the only (-1) -vertex. In this case, the chain Δ_1 is conjugate to both Δ_2 and $\Delta_3 - \diamond - \Delta_2$ (see Lemma 2.3.1), a contradiction. $\square$

6.5. Lemma. *Let $Q \in H \setminus \{P\}$ be any point and let $Q' \in \nu^{-1}(Q)$. Then $4 \geq \text{emb dim}(H, Q) \geq \text{emb dim}(H', Q') - 1$.*

Proof. By Corollary 6.2.1 the conductor ideal coincides with the ideal sheaf $I_{C'}$. The natural map $\mathcal{O}_H \rightarrow \nu_* \mathcal{O}_{H'}$ induces an isomorphism $I_C \simeq \nu_* I_{C'}$ (any regular function on H' that vanishes on C' descends to H). From the following commutative

digram

$$\begin{array}{ccccccc}
0 & \longrightarrow & \nu_* I_{C'} & \longrightarrow & \nu_* \mathcal{O}_{H'} & \longrightarrow & \nu_* \mathcal{O}_{C'} \longrightarrow 0 \\
& & \parallel & & \uparrow & & \uparrow \\
0 & \longrightarrow & I_C & \longrightarrow & \mathcal{O}_H & \longrightarrow & \mathcal{O}_C \longrightarrow 0
\end{array}$$

we have $\nu_* \mathcal{O}_{H'} / \mathcal{O}_H \simeq \nu_* \mathcal{O}_{C'} / \mathcal{O}_C$. Note that $\nu_* \mathcal{O}_{C'}$ is a locally free $\mathcal{O}_C$ -module and there is a local splitting $\nu_* \mathcal{O}_{C'} = \mathcal{O}_C \oplus \mathcal{O}_C t$ for some $t \in \nu_* \mathcal{O}_{C'}$. Thus $\nu_* \mathcal{O}_{H'} / \mathcal{O}_H \simeq \mathcal{O}_C t$. Therefore, $\mathfrak{m}_{Q',H'} / \mathfrak{m}_{Q',H'}^2$ is generated by $1 + \dim \mathfrak{m}_{Q,H} / \mathfrak{m}_{Q,H}^2$ elements as an $\mathcal{O}_{Q,H}$ -module. $\square$

6.5.1. Corollary (cf. [Tzi05]). *The chain Δ_3 in 6.4.1 satisfies the inequality*

$$(6.5.2) \quad \text{emb dim}(H', Q') - 3 = \sum (c_i - 2) \leq 2.$$

The proof of this statement is contained in [Tzi05, Proof of Th. 5.6] which is rather computational and uses the classification of degenerate cusp singularities. Here is a much shorter proof.

Proof. By Lemma 6.5 we have $\text{emb dim}(H', Q') \leq \text{emb dim}(H, Q) + 1 \leq 5$. On the other hand, since (H', Q') is a cyclic quotient singularity,

$$\begin{aligned}
\text{emb dim}(H', Q') &= -\left(\sum E_i\right)^2 + 1 = \\
&= 1 + \sum c_i - 2 \sum_{i \neq j} E_i \cdot E_j = 3 + \sum (c_i - 2),
\end{aligned}$$

where the E_i 's are exceptional divisors on the minimal resolution. This immediately gives the desired inequality. $\square$

6.6. Proposition. *Assume that we are in the case 6.4.1 under 6.1. Furthermore, assume that every member of $|\mathcal{O}_X|_C$ is non-normal and $\sum (c_i - 2) = 2$ (whence $\text{emb dim}(H, Q) = 4$). Let G (resp. G') be the fundamental cycle of $\Delta(H', C')$ (resp. Δ_3). Then $G \geq 2G'$ if and only if $\text{emb dim}(M, Q) = 4$ for general $M \in |\mathcal{O}_X|_C$.*

Proof. We have an analytic isomorphism $(H', Q') \simeq \mathbb{C}_{u,v}^2 / \mu_n(1, q)$ for some n, q with $\gcd(n, q) = 1$. By Proposition 6.3 the graph $\tilde{C}_1 - \Delta_3 - \tilde{C}_2$ is contracted on $\overline{H}$. Note that G is $\overline{\nu}$ -numerically trivial. Thus there is a function $\psi \in \mathcal{O}_H$ such that $\mu^* \nu^* \psi = 0$ defines G near $\mu^{-1} \nu^{-1}(Q)$. Hence the lifting of $\nu^* \psi$ to $\mathbb{C}_{u,v}^2$ is given by an invariant monomial λ multiplied by a unit.

Since $\sum (c_i - 2) = 2$, we see $\text{emb dim}(H', Q') = 5$ and $\text{emb dim}(H, Q) = 4$ by Corollary 6.5.1 and Lemma 6.5, and $I_{C'} \subset \mathfrak{m}'_{Q',H'}$ is generated by exactly three invariant monomials in u, v divisible by uv . Thus every minimal generating set of $I_C \subset \mathfrak{m}_{Q,H}$ induces a minimal generating set of $I_{C'} \subset \mathfrak{m}'_{Q',H'}$ (cf. the proof of Lemma 6.5). This means that $\text{emb dim}(M, Q) < 4$ for general $M \in |\mathcal{O}_X|_C$ iff $\nu^* \psi$ can be a part of a coordinate of (H', Q') . However since the lifting of $\nu^* \psi$ is an invariant monomial (times a unit), this happens iff $\nu^* \psi$ equals one of the three monomial generators of $I_{C'}$.

There are only two series of possibilities for $\Delta(H, C)$ near Q :

$$(*) \quad \diamond - \underbrace{\circ \cdots \circ}_{a-2} - \overset{4}{\circ} - \underbrace{\circ \cdots \circ}_{b-2} - \diamond \quad a, b \geq 2$$

$$(**) \quad \diamond - \underbrace{\circ \cdots \circ}_{a-2} - \overset{3}{\circ} - \underbrace{\circ \cdots \circ}_{b-2} - \overset{3}{\circ} - \underbrace{\circ \cdots \circ}_{c-2} - \diamond \quad a, b, c \geq 2$$

Each monomial in $I_{C'}$ corresponds to an effective divisor of $\tilde{H}$ with support $\Delta_3 \cup \tilde{C}$ which is μ -trivial (i.e., numerically trivial along Δ_3). The following table gives three such monomials (or divisors) m_A, m_B, m_C for each of (*) and (**). For instance the numbers of the row m_A shows the coefficient of the curve corresponding to the vertex in the divisor m_A .

(*)	$\diamond$	$\circ \dots\dots \circ$	$\overset{4}{\circ}$	$\circ \dots\dots \circ$	$\diamond$		
m_A	1	$\dots\dots\dots 1$	1	3 $\dots\dots\dots$	$2b-1$		
m_B	$2a-1$	$\dots\dots\dots 3$	1	1 $\dots\dots\dots$	1		
m_C	a	$\dots\dots\dots 2$	1	2 $\dots\dots\dots$	b		
(**)	$\diamond$	$\circ \dots \circ$	$\overset{3}{\circ}$	$\circ \dots \circ$	$\overset{3}{\circ}$	$\circ \dots \circ$	$\diamond$
m_A	1	$\dots\dots\dots$	1	$\dots\dots\dots$	b	$\dots\dots\dots$	$bc+c-1$
m_B	$ab+a-1$	$\dots\dots\dots$	b	$\dots\dots\dots$	1	$\dots\dots\dots$	1
m_C	a	$\dots\dots\dots$	1	$\dots\dots\dots$	1	$\dots\dots\dots$	c

It is clear that none of these monomials belong to $\mathfrak{m}_{Q',H'}^2$ because each vanishes to order 1 at one of the vertices with weight 3 or 4. Hence m_A, m_B, m_C are the monomial generators of $I_{Q'}$. One can also check that the lifting of $\nu^*\psi$ equals one of m_A, m_B, m_C iff one of the vertices of weight 3 or 4 appears with coefficient 1 in G iff $G \not\geq 2G'$. $\square$

6.7. Proposition. *Assume that f is a $\mathbb{Q}$ -conic bundle germ such that every member of $|\mathcal{O}_X|_C$ is non-normal. Assume furthermore that $H \in |\mathcal{O}_X|_C$ is taken to be general. Then C' is irreducible.*

6.7.1. Remark. If in the above assumptions X is of index 2, then $\Delta(H', C' + \Upsilon)$ is of the form

$$\square - \circ - \bullet - \circ - \square.$$

Proof of 6.7. Assume that C' is reducible. Then the dual graph $\Delta(H', C')$ is of the form 6.4.1 with $\diamond^2 = -1$. Clearly the chains Δ_1 and Δ_2 are not empty (otherwise X is Gorenstein). Since the matrix corresponding to $\bullet - \Delta_3 - \bullet$ is negative definite, the subgraph Δ_3 is not Du Val. We will use the inequality (6.5.2).

6.7.2. Assume that $r = s = 1$. Then $a_1 = b_1 = 2$ and the graph 6.4.1 or 6.4.2 is of the form

$$\circ - \bullet - \overset{4}{\circ} - \bullet - \circ \quad \text{or} \quad \circ - \bullet - \overset{3}{\circ} - \underbrace{\circ \cdots \circ}_l - \overset{3}{\circ} - \bullet - \circ \quad l \geq 0.$$

The fundamental cycle G of $\Delta(H', C')$ is given by

$$\overset{\circ}{1} - \overset{\bullet}{2} - \overset{\circ}{1} - \overset{\bullet}{2} - \overset{\circ}{1} \quad \text{or} \quad \overset{\circ}{1} - \overset{\bullet}{2} - \overset{\circ}{1} - \overset{\circ}{1} \cdots \overset{\circ}{1} - \overset{\circ}{1} - \overset{\bullet}{2} - \overset{\circ}{1},$$

respectively. Then by Proposition 6.6 our H is not general enough, a contradiction.

From now on we assume that $rs > 1$. Since $[a_1, \dots, a_r]$ and $[b_1, \dots, b_s]$ are conjugate, we may assume by symmetry that $a_1 = 2$, $b_1 > 2$, and $r > 1$.

6.7.3. Consider the case where the chain Δ_3 contains exactly one curve with self-intersection < -2 . Then the graph 6.4.1 has the following form:

$$\begin{array}{ccccccc} a_r & \cdots & a_1=2 & \bullet & \underbrace{\circ \cdots \circ}_{l_1} & \circ & \underbrace{\circ \cdots \circ}_{l_2} & \bullet & b_1 & \cdots & b_s \\ \circ & & \circ & & & & & & \circ & & \circ \end{array}$$

where $c = 3$ or 4 . Since $a_1 = 2$, it holds $l_1 = 0$ because the graph $\circ - \bullet - \circ$ is not negative definite. Choose the above configuration so that c is minimal.

If $l_2 > 0$, then contracting both black vertices we get

$$\begin{array}{ccccccc} a_r & \cdots & a_2 & \bullet & \underbrace{\circ \cdots \circ}_{l_2-1} & \bullet & b_1-1 & \cdots & b_s \\ \circ & & \circ & & & & \circ & & \circ \end{array}$$

The strings $[a_2, \dots, a_r]$ and $[b_1 - 1, \dots, b_s]$ at the ends are again conjugate. This contradicts our minimality assumption because $c' = c - 1 < 4$.

Therefore, $l_1 = l_2 = 0$ and 6.4.1 is of the form

$$\begin{array}{ccccccc} a_r & \cdots & a_1 & \bullet & \circ & \bullet & b_1 & \cdots & b_s \\ \circ & & \circ & & & & \circ & & \circ \end{array}$$

Contracting black vertices we get

$$\begin{array}{ccccccc} a_r & \cdots & a_2 & \bullet & \circ & \bullet & b_1-1 & \cdots & b_s \\ \circ & & \circ & & & & \circ & & \circ \end{array}$$

Hence $c = 4$ and $a_2 \geq 3$. Again the string $[a_2, \dots, a_r]$ is conjugate to both $[b_1 - 1, \dots, b_s]$ and $[c - 2, b_1 - 1, \dots, b_s]$, a contradiction.

6.7.4. Now we consider the case where Δ_3 contains exactly two (-3) -curves. Then the graph 6.4.1 has the following form:

$$\begin{array}{ccccccc} a_r & \cdots & a_1=2 & \bullet & \underbrace{\circ \cdots \circ}_{l_1} & \circ & \underbrace{\circ \cdots \circ}_{l_2} & \bullet & b_1 & \cdots & b_s \\ \circ & & \circ & & & & & & \circ & & \circ \end{array}$$

(As above $c_1 > 2$, since $a_1 = 2$). If $l_2 > 0$, then contracting both black vertices we get

$$\begin{array}{ccccccc} a_r & \cdots & a_2 & \bullet & \underbrace{\circ \cdots \circ}_{l_1+1} & \circ & \underbrace{\circ \cdots \circ}_{l_2-1} & \bullet & b_1-1 & \cdots & b_s \\ \circ & & \circ & & & & & & \circ & & \circ \end{array}$$

Here again the strings $[a_2, \dots, a_r]$ and $[b_1 - 1, \dots, b_s]$ are conjugate. This contradicts the case considered above. So, $l_2 = 0$. Then contracting both black vertices we get

$$\begin{array}{ccccccc} a_r & \cdots & a_2 & \bullet & \underbrace{\circ \cdots \circ}_{l_1+2} & \bullet & b_1-1 & \cdots & b_s \\ \circ & & \circ & & & & \circ & & \circ \end{array}$$

As above the string $[a_2, \dots, a_r]$ is conjugate to both $[b_1 - 1, \dots, b_s]$ and $[2, \dots, 2, b_1 - 1, \dots, b_s]$, a contradiction.

□

6.8. Corollary. Let f be a $\mathbb{Q}$ -conic bundle such that a general member $H \in |\mathcal{O}_X|_C$ is not normal. Then the germ (H, C) is analytically isomorphic to the germ along the line $L := \{y = z = 0\}$ of the hypersurface given by the following weighted polynomial of degree $2m$ in variables x, y, z, u :

$$\phi := x^{2m-2a}y^2 + x^{2a}z^2 + yzu.$$

in $\mathbb{P}(1, a, m-a, m)$, for some integers a, m such that $0 < a < m$ and $\gcd(a, m) = 1$.

Proof. By Proposition 6.7 (H, C) is of type 6.4.2. Then it is easy to see that the pair (H, C) up to analytic isomorphism is uniquely defined by the types of singularities $\frac{1}{m}(1, a)$ and $\frac{1}{m}(1, -a)$. On the other hand, the hypersurface $\phi = 0$ satisfies the conditions of 6.4.2. $\square$

Note that we are interested only in the *germ* of the hypersurface $\{\phi = 0\}$ along L .

6.8.1. Remark. Since the germ $(\{\phi = 0\}, L)$ is analytically isomorphic to our (H, C) , there is a rational curve fibration on $(\{\phi = 0\}, L)$ whose central fiber is L . One can check that this fibration is given by the rational function

$$s = \frac{y^{m-a}z^a}{x^{2a(m-a)}},$$

which is regular in a neighborhood of L in H .

6.8.2. Lemma. *Let (H, C) be as in 6.8 and let $s : H \rightarrow T$ be the corresponding rational curve fibration. Let $t : X \rightarrow \mathbb{C}$ be a one-parameter smoothing of (H, C) in a $\mathbb{Q}$ -Gorenstein family. If X has only terminal singularities, then (X, C) is a $\mathbb{Q}$ -conic bundle germ.*

Proof. Let $V := s^{-1}(o)$ (with the scheme structure) and let Z be the component of the Hilbert scheme of X containing the point $o = [V]$ representing V . Let $\mathfrak{X} \subset X \times Z$ be the corresponding universal family. We have the following commutative diagram

$$\begin{array}{ccc} & V & \xlongequal{\quad} W \\ & \swarrow & \searrow \\ X & \xleftarrow{p} & \mathfrak{X} \\ \downarrow t & \swarrow \text{dotted} & \downarrow \pi \\ \mathbb{C} & \xleftarrow{\quad} & Z \end{array}$$

where $W := \pi^{-1}(o)$. Both V and W are locally complete intersections. Moreover, $I_V/I_V^2 \simeq \mathcal{O}_V \oplus \mathcal{O}_V$ and $I_W/I_W^2 \simeq \mathcal{O}_W \oplus \mathcal{O}_W$. Since $H^1(V, (I_V/I_V^2)^\vee) = 0$, Z is smooth at o and there is a natural isomorphism $\mathbb{C}^2 \simeq T_{o,Z} \simeq H^0(V, (I_V/I_V^2)^\vee)$. On the other hand, $H^0(V, (I_W/I_W^2)^\vee) \simeq T_{o,Z}$ because W is a fiber of π . Therefore, there is a natural isomorphism $H^0(W, (I_W/I_W^2)^\vee) \simeq H^0(V, (I_V/I_V^2)^\vee)$ and the natural map $(I_W/I_W^2)^\vee \rightarrow (I_V/I_V^2)^\vee$ is also an isomorphism. Thus p is an isomorphism in a neighborhood of W . By shrinking $\mathfrak{X}$ and X we may assume that there is a contraction $X \rightarrow Z$ such that the whole diagram is commutative. $\square$

The existence of a $\mathbb{Q}$ -Gorenstein smoothing follows from [Tzi09]. However in our particular case we can construct it explicitly:

6.8.3. Lemma. *Let (H, C) , m, a be as in Corollary 6.8. For $s = (s_1, \dots, s_5) \in \mathbb{C}_s^5$, hypersurfaces $H_s \subset \mathbb{P}(1, a, m-a, m)$ given by equation*

$$\phi_s := \phi + s_1 x^{2m-a} y + s_2 x^{m-a} u y + s_3 x^{2m} + s_4 x^m u + s_5 u^2 = 0$$

form a miniversal deformation family of the germ $C \subset H$.

Proof. We compute $T_{qG}^1(H)$ from the $\mathbb{Q}$ -Gorenstein smoothing $H \subset P := \mathbb{P}(1, a, m - a, m)$ (cf. [Tzi09, §3]). By definition, $T_{qG}^1(H)$ has an ℓ -structure and $T_{qG}^1(H)^\# = T_{qG}^1(H^\#)$. Furthermore, we get an exact sequence

$$\mathcal{H}om_H(\Omega_P^1, \mathcal{O}_H) \longrightarrow \mathcal{H}om_H(\mathcal{O}_P(-H), \mathcal{O}_H) \longrightarrow T_{qG}^1(H) \rightarrow 0$$

of sheaves with ℓ -structures. So $T_{qG}^1(H) = \mathcal{O}_P(2m)/G$, where G is generated by ϕ and its derivatives. A direct computation shows that $x^{2m-a}y$, $x^{m-a}yu$, x^{2m} , $x^m u$, u^2 form a $\mathbb{C}$ -basis of the vector space $T_{qG}^1(H)$; $x^{2m-a}y$, $x^{m-a}yu$ generate the torsion part of $T_{qG}^1(H)$ and x^{2m} , $x^m u$, u^2 generate $T_{qG}^1(H)/(\text{torsion}) \simeq \mathcal{O}_P(2m) \otimes \mathcal{O}_C \simeq \mathcal{O}_{\mathbb{P}^1}(2)$. $\square$

6.8.4. Example. Let $\alpha, \beta \in \mathbb{C}$ are some general constant and let X be the threefold given in $\mathbb{P}(1, a, m - a, m) \times \mathbb{C}_t$ by

$$\phi + (\alpha x^m - u)(\beta x^m - u)t = 0.$$

Then the singularities of X along the curve $C := \{y = z = t = 0\}$ consists of a cyclic quotient singularity of type $\frac{1}{m}(1, a, m - a)$ at $\{x = y = z = t = 0\}$ and two (Gorenstein) ordinary double points at $\{\alpha x^m - u = y = z = t = 0\}$ and $\{\beta x^m - u = y = z = t = 0\}$. The contraction $X \rightarrow Z$ exists by Lemma 6.8.2.

Thus Theorem 1.5 is proved. Now assume that f is birational.

6.9. Lemma ([Tzi05, Th. 5.6, (1a)]). *If f is birational, then C' is reducible and the dual graph $\Delta(H', C')$ is of the form 6.4.1.*

Proof. Assume that $\Delta(H', C')$ is of the form 6.4.2. Then the chain of smooth rational curves corresponding to the graph $\Delta_1 - \bullet - \Delta_2$ is contracted by v . On the other hand, Δ_1 and Δ_2 are conjugate. By Lemma 2.3.1 this configuration corresponds to a rational curve fibration, i.e., v is not birational, a contradiction. $\square$

6.10. The singularity (H, Q) is a so-called *degenerate cusp* [SB83]. One can define the *fundamental cycle* Γ of (H, Q) and attach an invariant $\zeta = -\Gamma^2$ to (H, Q) such that

$$\begin{aligned} \zeta = 1 &\iff (H', Q') \text{ is a smooth point} \iff (H, Q) \simeq \{y^2 = x^3 + x^2 z^2\}, \\ \zeta = 2 &\iff (H', Q') \text{ is a Du Val point} \iff (H, Q) \simeq \{y^2 = x^2 z^2 + x^{n+3}\}, \\ &\quad \text{of type } A_n, n \geq 1 \\ \zeta = 3 &\iff \sum (c_i - 2) = 1 \iff (H, Q) \simeq \{xyz = y^{a+3} + z^{b+3}\}, \\ &\quad a, b \geq 0, \\ \zeta = 4 &\iff \sum (c_i - 2) = 2 \iff \text{emb dim}(H, Q) = 4, \end{aligned}$$

(see [SB83, §1]). Then by [Tzi09, Th. 3.1, Prop. 3.4] we have

6.10.1. Theorem. *In the above notations, a one parameter smoothing of (H, C) with only terminal singularities exists if and only if*

$$\tilde{C}_1^2 + \tilde{C}_2^2 + 1 + 4\delta_{\zeta,1} + 4\delta_{\zeta,2} + 3\delta_{\zeta,3} + 2\delta_{\zeta,4} \geq 0,$$

where $\delta_{i,j}$ is Kronecker's delta.

6.10.2. Remark. One can see that the last inequality is equivalent to

$$\tilde{C}_1^2 + \tilde{C}_2^2 + 5 - \sum (c_i - 2) \geq 0,$$

where we put $\sum (c_i - 2) = 0$ if Δ_3 is empty.

6.10.3. Example. Assume that the configuration in 6.4.1 is of the form

$$\circ \text{---} \overset{4}{\circ} \text{---} \overset{4}{\diamond} \text{---} \bullet \text{---} \circ \text{---} \circ \text{---} \overset{3}{\circ}$$

Then (X, C) is a divisorial extremal neighborhood. By Proposition 6.3 every member of $|\mathcal{O}_X|_C$ is non-normal. By 6.10 this H is general in $|\mathcal{O}_X|_C$.

7. ON INDEX TWO $\mathbb{Q}$ -CONIC BUNDLES

In this section we give examples of index two $\mathbb{Q}$ -conic bundles. Let $y_1, y_2, y_3, y_4; u, v$ be as in Theorem 1.6, and let $X \subset \mathbb{P}(1, 1, 1, 2) \times \mathbb{C}^2$ given by

$$\begin{aligned} 0 &= \alpha_1 y_1^2 + \alpha_2 u^e y_4 + (\beta_2 u + v) y_3^2 \\ 0 &= \alpha_3 (y_2^2 + \beta_1 y_1 y_3) + \alpha_4 u y_3^2 + v y_4, \end{aligned}$$

where $\alpha_1, \dots, \alpha_4 \in \mathbb{C}$ are general, $\beta_1, \beta_2 \in \mathbb{C}$ are either zero or general, and $e = 1, 2, 3$. Furthermore, $C \subset X$ is given by $y_1 = y_2 = u = v = 0$.

By Bertini's Theorem, we see that the singular locus, Σ , of X is contained in $\{u = v = 0\}$. Hence, $\Sigma \subset \{u = v = y_1 = y_2 = 0\}$ and using notation $[y : z] := (0 : 0 : y : z) \times (0, 0)$, we see

$$\begin{aligned} \Sigma &= \left\{ [y_3 : y_4] \mid \text{rank} \begin{pmatrix} 0 & \alpha_2 e u^{e-1} y_4 + \beta_2 y_3^2 & y_3^2 \\ \beta_1 y_3 & \alpha_4 y_3^2 & y_4 \end{pmatrix} \leq 1 \right\} \cup \{[0 : 1]\} \\ &= \begin{cases} \{[0 : 1]\} & \text{if } \beta_1 \neq 0, \\ \left\{ \left[1 : \pm \sqrt{\alpha_4 / \alpha_2} \right], [0 : 1] \right\} & \text{if } \beta_1 = 0, \beta_2 = 0, \text{ and } e = 1, \\ \{[0 : 1]\} & \text{if } \beta_1 = 0, \beta_2 = 0, \text{ and } e > 1, \\ \{[1 : \alpha_4 / \beta_2], [0 : 1]\} & \text{if } \beta_1 = 0, \beta_2 \neq 0, \text{ and } e > 1. \end{cases} \end{aligned}$$

At $[0 : 1]$, the singularity $(X, [0 : 1])$ is a hyper-quotient:

$$\{\alpha_1 y_1^2 + \alpha_2 u^e + \beta_2 u y_3^2 - \alpha_3 y_2^2 y_3^2 - \alpha_3 \beta_1 y_1 y_3^3 - \alpha_4 u y_3^4 = 0\} / \mu_2(1, 1, 1, 0).$$

By [Mor85, Cor. 2.1], we see that $(X, [0 : 1])$ is a terminal singularity of type

- $\frac{1}{2}(1, 1, 1)$ if $e = 1$,
- cAx/2 if $e = 2$ (cf. [Mor85, Thm. 12, (3)]),
- cD/2 if $e = 3$ and $\beta_2 \neq 0$ [Mor85, Thm. 23],
- cE/2 if $e = 3$ and $\beta_2 = 0$ (cf. [Mor85, Thm. 25]).

Every other singular point, if any, is easily seen to be an ordinary double point, in particular a type (III) point.

- (i) Case $\beta_1 \neq 0$: In this case we can assume $\beta_1 = -1$ by change of coordinate $y_1 \mapsto -y_1/\beta_1$, and we are in case (i) of Theorem 1.6. In this case, $[0 : 1]$ is the only singular point and it can be of type $\frac{1}{2}(1, 1, 1)$, cAx/2, cD/2 or cE/2 as above.
- (ii) Case $\beta_1 = 0$: In this case we are in case (ii) of Theorem 1.6. The type of singularity of (X, C) in our example is
 - $\frac{1}{2}(1, 1, 1) + (\text{III}) + (\text{III})$ if $\beta_2 = 0$ and $e = 1$,

- $cAx/2 + (III)$ if $\beta_2 \neq 0$ and $e = 2$,
- $cAx/2$ if $\beta_2 = 0$ and $e = 2$,
- $cD/2 + (III)$ if $\beta_2 \neq 0$ and $e = 3$, and
- $cE/2$ if $\beta = 0$ and $e = 3$.

In particular, we have shown that all types of terminal index two singularities can appear on $\mathbb{Q}$ -conic bundles as in Theorem 1.6.

Acknowledgements. The authors are grateful to Professors A. Fujiki and O. Fujino for helpful discussions. The paper was written when the second author was visiting Research Institute of Mathematical Sciences and Mathematical Department of Kyoto University in 2009. He would like to thank both institutions for the hospitality and excellent working environment. The first author was partially supported by JSPS Grant-in-Aid for Scientific Research (B)(2), No. 20340005. The second author was partially supported by grants RFBR, No. 08-01-00395-a and Kyoto University's Global COE.

REFERENCES

- [Art68] M. Artin, On the solutions of analytic equations. *Invent. math.*, 5: 277–291 (1968).
- [Bin81] J. Bingener. On the existence of analytic contractions. *Invent. Math.*, 64(1): 25–67, 1981.
- [KM92] J. Kollár and S. Mori. Classification of three-dimensional flips. *J. Amer. Math. Soc.*, 5(3):533–703, 1992.
- [Kol92] J. Kollár, editor. *Flips and abundance for algebraic threefolds*. Société Mathématique de France, Paris, 1992. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991, Astérisque No. **211** (1992).
- [Fuj79] A. Fujiki. Closedness of the Douady spaces of compact Kähler spaces. *Publ. Res. Inst. Math. Sci.*, 14(1): 1–52, 1978/79.
- [Gro68] A. Grothendieck. *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*. North-Holland Publishing Co., Amsterdam, 1968. Augmenté d'un exposé par Michèle Raynaud, Séminaire de Géométrie Algébrique du Bois-Marie, 1962, Advanced Studies in Pure Mathematics, Vol. 2.
- [Kaw88] Y. Kawamata. Crepant blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces. *Ann. of Math. (2)*, 127(1):93–163, 1988.
- [KSB88] J. Kollár and N. I. Shepherd-Barron. Threefolds and deformations of surface singularities. *Invent. Math.*, 91(2):299–338, 1988.
- [LW86] E. Looijenga and J. Wahl. Quadratic functions and smoothing surface singularities. *Topology*, 25(3):261–291, 1986.
- [Mor85] S. Mori. On 3-dimensional terminal singularities. *Nagoya Math. J.*, 98:43–66, 1985.
- [Mor88] S. Mori. Flip theorem and the existence of minimal models for 3-folds. *J. Amer. Math. Soc.*, 1(1):117–253, 1988.
- [Mor02] S. Mori. On semistable extremal neighborhoods. In *Higher dimensional birational geometry (Kyoto, 1997)*, *Adv. Stud. Pure Math.* **35**, 157–184. Math. Soc. Japan, Tokyo, 2002.
- [MP08] S. Mori and Y. Prokhorov. On $\mathbb{Q}$ -conic bundles. *Publ. Res. Inst. Math. Sci.*, 44(2):315–369, 2008.
- [MP09] S. Mori and Y. Prokhorov. On $\mathbb{Q}$ -conic bundles, III. *Publ. Res. Inst. Math. Sci.*, 45:787–810, 2009.
- [Nak87] N. Nakayama. The lower semicontinuity of the plurigena of complex varieties. In *Algebraic geometry, Sendai, 1985*, *Adv. Stud. Pure Math.* **10**, 551–590. North-Holland, Amsterdam, 1987.
- [Pro97] Y. Prokhorov. On the complementability of the canonical divisor for Mori fibrations on conics. *Sbornik. Math.*, 188(11):1665–1685, 1997.
- [Pro01] Y. Prokhorov. *Lectures on complements on log surfaces*, *MSJ Memoirs* **10**. Mathematical Society of Japan, Tokyo, 2001.

- [Rei83] M. Reid. Minimal models of canonical 3-folds. In *Algebraic varieties and analytic varieties (Tokyo, 1981)*, *Adv. Stud. Pure Math.* **1**, 131–180. North-Holland, Amsterdam, 1983.
- [Rei87] M. Reid. Young person’s guide to canonical singularities. In *Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985)*, *Proc. Sympos. Pure Math.* **46**, 345–414. Amer. Math. Soc., Providence, RI, 1987.
- [SB83] N. I. Shepherd-Barron. Degenerations with numerically effective canonical divisor. In *The birational geometry of degenerations (Cambridge, Mass., 1981)*, *Progr. Math.* **29**, 33–84. Birkhäuser Boston, Boston, MA, 1983.
- [Sho93] V. V. Shokurov. 3-fold log flips. *Russ. Acad. Sci., Izv., Math.*, 40(1):95–202, 1993.
- [Ste88] J. Stevens. On canonical singularities as total spaces of deformations. *Abh. Math. Sem. Univ. Hamburg*, 58:275–283, 1988.
- [Tzi05] N. Tziolas. Three dimensional divisorial extremal neighborhoods. *Math. Ann.*, 333(2):315–354, 2005.
- [Tzi09] N. Tziolas. Q-Gorenstein deformations of nonnormal surfaces. *Amer. J. Math.*, 131(1):171–193, 2009.

RIMS, KYOTO UNIVERSITY, OIWAKE-CHO, KITASHIRAKAWA, SAKYO-KU, KYOTO 606-8502, JAPAN

E-mail address: `mori@kurims.kyoto-u.ac.jp`

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS, MOSCOW STATE UNIVERSITY, MOSCOW 117234, RUSSIA

E-mail address: `prokhorogmail.com`